

cases (13/42) of these: 4 cases of 45, XO, 4 cases of trisomy 18, 4 cases of trisomy 21, and 1 case of trisomy 13. Among fetuses with normal karyotype (n= 29), intrauterine fetal death occurred in two fetuses before 20 weeks of gestation. Major structural fetal malformations detected in 20-weeks level II sonogram were diagnosed in 11 of the remaining 27 cases (40.7%). A total of 29 chromosomally normal pregnancies gave rise to 16 live births (55.2%), with no structural abnormalities.

CONCLUSION: These data suggest that the prognosis of fetus with CH diagnosed in the first trimester is poor. Therefore, fetuses with CH requires more careful assessment of the fetus, with regard to both karyotyping and follow-up level II scan for detecting structural anomaly.

0002-9378/\$ – see front matter • doi:10.1016/j.ajog.2009.10.396

382 Customized vs population-based approach to evaluate fetal overgrowth

Maged Costantine¹

¹for the Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network, Bethesda, Maryland

OBJECTIVE: To evaluate a customized method to estimate fetal overgrowth in mild gestational diabetes mellitus (mGDM) and normal pregnancies

STUDY DESIGN: Secondary analysis of a study with 4 cohorts: treated mGDM, untreated mGDM, positive GLT screen but normal OGTT, and normal. The 1° neonatal outcome was a composite of hypoglycemia, hyperbilirubinemia, hyperinsulinemia, trauma, or perinatal death. LGA>90th%ile was evaluated using ethnicity & gender-specific population (pop) norms (Alexander), and customized (cust) birth weight (BW) centile based also on maternal height, weight, and parity (Gardosi et al). LGAcust and LGApop were compared for their association with outcome. LGAcust was correlated with neonatal fat mass.

RESULTS: 642 (32.1%) infants developed the 1° outcome. Mean custBW centile was higher in those who developed the 1° outcome compared with those who did not (59.5±29.6 vs. 50.0±29.3, p<0.0001). Of those who had the 1° outcome 19.3% and 13.2% were LGAcust and LGApop, vs. 11.3% and 8.2% for those who did not (difference 6.1 vs. 3.1%, P=.03). Among normal women, 9.9% were LGAcust, close to expected 10%, vs. 6.8% LGA pop (p=0.03). Cust BW centile was also significantly correlated with neonatal fat mass (r=0.70, p<.0001).

CONCLUSION: A customized approach for assessment of fetal growth, instead of current population-based norms, better identifies neonates at risk for adverse outcomes related to overgrowth.

Cohort (N)	Norm	% LGA with 1° outcome	% LGA without 1° outcome
treated GDM (422)	pop	8.0	6.0
	cust	12.4	11.2
Untreated GDM (400)	pop	19.7	11.9
	cust	32.7	16.6
Pos GLT (798)	pop	13.1	9.1
	cust	15.8	11.0
Normal (381)	pop	11.2	5.7
	cust	18.4	7.4

0002-9378/\$ – see front matter • doi:10.1016/j.ajog.2009.10.397

383 Physiological basis of the Quintero staging system

Luminita Crisan¹, Eftichia Kontopoulos², Ramen Chmait³, Ruben Quintero²

¹University of South Florida, Tampa, Florida, ²University of Miami, Miami, Florida, ³University of Southern California, Los Angeles, California

OBJECTIVE: Twin-twin transfusion syndrome (TTTS) is thought to result from a net flow of blood from the donor to the recipient twin. The Quintero staging system (QSS) is used to classify the heterogeneity and gauge the severity of TTTS and assumes increased severity with advancing Stage. The purpose of this study was to assess the relationship between the flow in the umbilical vein (UV) of the donor twin and Stage.

STUDY DESIGN: UV flow (cc/min) was calculated as: $Q_{uv} = A_u \times V_u \times 60$, where A_u and V_u are the area and the blood flow velocity of the UV, respectively. The diameter of the UV was obtained by ultrasound in a perpendicular view of the vein, where as the velocity was obtained in a parallel view. Pulsatile V_u was assessed as the time-average velocity by trace. Measurements were obtained prior to laser surgery in all cases. Logistic regression controlled for Stage, gestational age and estimated fetal weight. Surgeries were approved by the Institutional Review Board and all patients signed informed consent.

RESULTS: Three hundred and thirty-two patients underwent pre-operative assessment. Table I shows the median UV flow values by QSS. There was a significant inverse correlation between UV flow and QSS, with a decrease in UV flow from Stage I-III and increased flow in Stage IV. The estimated fetal weight was also significantly correlated, but did not differentiate Stages I-II.

CONCLUSION: Umbilical venous flow assessment in the donor twin correlates inversely with disease severity as gauged by the QSS from Stages I-III, differentiating one Stage from another. Improved blood flow to the donor in Stage IV may be associated with increased venous pressure in the recipient twin from congestive heart failure. These hemodynamic findings support the basis of the Quintero Staging system.

Table I. Pre-operative umbilical venous flow in the donor twin by Quintero Stage

Stage (N)	UV flow (cc/min)	P value
I (64)	65	
II (95)	51	.007
III (149)	32	.0001
IV (24)	80	.0001

0002-9378/\$ – see front matter • doi:10.1016/j.ajog.2009.10.398

384 Does fetal echocardiogram perform better than standard fetal ultrasound in obese women?

Meredith Cruz¹, Maria Colon¹, Isabelle Wilkins²

¹University of Illinois at Chicago, Chicago, Illinois,

²University of Illinois at Chicago, Illinois

OBJECTIVE: Only 20-35% of major congenital cardiac anomalies are detected by routine ultrasound screening. It is often suggested that patients at risk should therefore undergo a fetal echocardiogram. We examined the accuracy of fetal echocardiogram when compared to standard fetal ultrasound in detecting cardiac anomalies, stratified by maternal BMI.

STUDY DESIGN: Retrospective chart review. Neonates with discharge diagnosis of any congenital cardiac anomaly between 2003 and 2008 were included. Maternal charts were reviewed for gestational age at ultrasound, ultrasound diagnosis, echocardiogram diagnosis, and maternal BMI. Subjects were classified into BMI categories by CDC criteria: underweight and normal (control): 18.5-24.9 kg/m², overweight: 25.0-29.9 kg/m², and obese: 30 kg/m² and above. Detection rates between the two modalities were calculated and compared. Accuracy of the ultrasound and echocardiogram diagnoses defined as