

**21** Integrase strand transfer inhibitors given to HIV-infected women late in pregnancy decrease HIV viral load more quickly than other antiretroviral therapy (ART)

Lisa Rahangdale<sup>1</sup>, Jordan Cates<sup>1</sup>, JoNell Potter<sup>2</sup>, Martina L. Badell<sup>3</sup>, Dominika Seidman<sup>4</sup>, Emily S. Miller<sup>5</sup>, Jenell S. Coleman<sup>6</sup>, Gweneth B. Lazenby<sup>7</sup>, Judy Levison<sup>8</sup>, William R. Short<sup>9</sup>, Sigal Yawetz<sup>10</sup>, Andrea Ciaranello<sup>11</sup>, Elizabeth Livingston<sup>12</sup>, Lunthita Duthely<sup>2</sup>, Bassam H. Rimawi<sup>13</sup>, Jean R. Anderson<sup>6</sup>, Elizabeth Stringer<sup>1</sup>

<sup>1</sup>University of North Carolina, Chapel Hill, NC, <sup>2</sup>University of Miami, Miami, FL, <sup>3</sup>Emory University, Atlanta, GA, <sup>4</sup>University of California San Francisco, San Francisco, CA, <sup>5</sup>Northwestern University Feinberg School of Medicine, Chicago, IL, <sup>6</sup>Johns Hopkins University, Baltimore, MD, <sup>7</sup>Medical University of South Carolina, Charleston, SC, <sup>8</sup>Baylor College of Medicine, Houston, TX, <sup>9</sup>University of Pennsylvania, Philadelphia, PA, <sup>10</sup>Brigham & Women's Hospital, Boston, MA, <sup>11</sup>Massachusetts General Hospital, Boston, MA, <sup>12</sup>Duke University, Durham, NC, <sup>13</sup>Emory University, Atlanta, GA

**OBJECTIVE:** Minimizing time to HIV viral suppression is critical in pregnancy. Integrase strand transfer inhibitors (INSTIs), like raltegravir, are known to rapidly suppress plasma HIV RNA in nonpregnant adults. We describe time to clinically relevant reduction in HIV RNA in pregnant women using INSTI-containing and non-INSTI-containing ART options.

**STUDY DESIGN:** We conducted a retrospective cohort study of pregnant HIV-infected women in the U.S. from 2009 to 2015. We included women who initiated ART, intensified their regimen or switched to a new regimen due to detectable viremia (HIV RNA > 40c/mL) at > 20 weeks gestation. Among women with a baseline HIV RNA permitting 1-log reduction, we estimated time to 1-log RNA reduction using the Kaplan-Meier estimator comparing women starting/adding an INSTI in their regimen versus other ART. To compare groups with similar follow-up time, we also conducted a subgroup analysis limited to women with <14 days between baseline and follow-up RNA data.


**RESULTS:** Maternal characteristics of 101 women from 11 U.S. clinics are shown in Table 1. 76/101 (75%) women were not taking ART at baseline. 39/101 (39%) women started/added an INSTI in their ART regimen. Among 90 women with a baseline HIV RNA permitting 1-log reduction, the median time to 1-log RNA reduction was 8 [IQR: 6, 14] days in the INSTI group versus 35 [IQR: 20, 53] days (p<0.001) in the non-INSTI ART group. In a subgroup of 39 women with first and last RNA measurements <14 days apart, median time to 1-log reduction was 7 [IQR: 6, 10] days in the INSTI group versus 11 [IQR: 10, 14] days (p=0.01) in the non-INSTI group. (Figure 1)

**CONCLUSION:** ART that includes INSTIs appears to induce more rapid viral suppression than other ART regimens in pregnancy. Inclusion of an INSTI may play a role in optimal reduction of HIV-RNA for HIV-infected pregnant women presenting late to care or failing initial therapy. Larger studies are urgently needed to assess the safety and effectiveness of this approach.

TABLE 1. Characteristics of 101 HIV-infected pregnant women from 11 US clinics post 20 weeks gestation at time of starting/including a new ART regimen, 2009-2015. Categorical variables are expressed as N (%), and continuous variables are expressed as median (IQR).

	Integrase (N=39)	Other ART (N=62)	p-value
Race & Ethnicity			
White, non-Hispanic	5 (13)	11 (18)	0.47
White, Hispanic	4 (11)	6 (10)	
Black	28 (72)	45 (73)	
Other	2 (5)	0 (0)	
Parity			
0	17 (44)	21 (34)	0.63
≥1	22 (56)	40 (65)	
Maternal Age (years)	29 (23, 34)	26 (23, 31)	0.33
Median Gestational Age (weeks)	33.6 (29.9-36.1)	27.5 (25.4-30.7)	<0.01
Body Mass Index (kg/m <sup>2</sup> )	28.2 (24.4-35.0)	30.5 (26.6-35.3)	0.33
No ART at baseline	20 (51)	56 (90)	<0.01
Initial HIV RNA (log <sub>10</sub> )	4.3 (3.5-4.9)	4.1 (3.3-4.6)	0.17
AIDS Diagnosis (copies/mL)	21,278 (3,370-71,660)	13,015 (2,050-35,570)	<0.01
	23 (60)	20 (32)	

FIGURE 1. Cumulative incidence of time to 1-log<sub>10</sub> change among A) 90 HIV-infected pregnant women from 11 U.S. clinics and B) a subset of 39 of the 90 HIV-infected pregnant women with baseline and follow up HIV RNA measurement of less than 14 days, stratified by drug regimen intervention.



**22** Changes in the patterns and rates of term stillbirth in the USA following the adoption of the 39-week rule: a cause for concern?

James Nicholson<sup>1</sup>, Lisa Kellar<sup>2</sup>, Shala Ahmad<sup>1</sup>, Ayesha Abid<sup>1</sup>, Jason Woloski<sup>1</sup>, Nadine Hewamudalige<sup>1</sup>, Serdar Ural<sup>1</sup>, Jerome Yaklic<sup>2</sup>

<sup>1</sup>Penn State Hershey Medical Center, Hershey, PA, <sup>2</sup>Boonshoft School of Medicine, Wright State University, Dayton, OH

**OBJECTIVE:** Because of reported associations between early-term delivery and various neonatal morbidities, an obstetric guideline called “the 39-week rule” was formally implemented in 2010 throughout the USA. The rule restricts “non-indicated” planned delivery prior to 39 weeks of gestation. However, several publications have suggested that the rule, by increasing the mean gestational age of term delivery, might increase population rates of term stillbirth. The objective of this large study was to determine if implementation of the 39-week rule was associated with an increase in the USA rate of term stillbirth.

**STUDY DESIGN:** Sequential ecological study, using state data, of the patterns and rates of term live births and term stillbirths within the USA between 2007 and 2013, inclusive.

**RESULTS:** Usable datasets were obtained from 46 states. Between 2007 and 2013 there was a continuous decrease in the proportion of term deliveries occurring in the 37th and 38th weeks of gestation. Over the same interval there was a continuous increase in the rate of term stillbirth (slope: 0.0186/1000/year, 95% CI 0.002-0.035). When comparing the 2007-2009 period to the 2011-2013 period, the rate of USA term stillbirth increased from 1.103/1000 to 1.177/1000 (RR 1.067, 95% CI 1.038-1.096).

**CONCLUSION:** Between 2007 and 2013 in the USA, the implementation of the 39-week rule achieved its primary goal of reducing the proportion of term births occurring before the 39th week of