

$p = 0.045$). The proportion of patients delivered at 200 minutes was 19% in the dextrose group versus 8% in the normal saline group. The proportion of patients delivered at 450 minutes was 75% in the dextrose group versus 61% in the normal saline group. There were no difference in the rate of cesarean section and APGAR score.

CONCLUSION: Glucose supplementation significantly reduces the length of the first and second stages of labour without increasing the rate of complication in induced, nulliparous women. Given the low-cost and the safety of this intervention, glucose should be used as the default solute during labour.

33 Randomized control trial of IV acetaminophen for post cesarean delivery pain control



Brie Altenau¹, Catrina Crisp¹, Ganga Devaiah², Donna Lambers¹

¹TriHealth, Cincinnati, OH, ²TriHealth - Hatton Research Institute, Cincinnati, OH

OBJECTIVE: To determine if administration of IV acetaminophen following routine scheduled cesarean section delivery would decrease the need for narcotic medications to control post-operative pain.

STUDY DESIGN: This was an IRB approved, double-blind, placebo-controlled randomized trial, registered on clinicaltrials.gov (NCT02046382). Women scheduled to undergo routine cesarean section for singleton pregnancy at term were recruited for enrollment. Study patients were given all medications in a standardized manner receiving either acetaminophen 1000mg intravenous (IV) every 8 hours for 48 hours or saline (placebo) for a total of 6 doses. The pharmacy prepared IV acetaminophen and saline in identical administration bags labeled “study drug” to ensure blinding. The initial dose of “study drug” was given within 60 minutes of skin incision. Quantity of breakthrough and scheduled analgesic medications, self-reported pain on the Faces Pain Scale (0-10) before and after study drug administration, and demographics were extracted from the patients’ charts. Power calculation determined that 45 patients per study arm were required to detect a 30% reduction in post-cesarean narcotic requirement with 80% power and a significance level of 0.05.

RESULTS: 133 patients were consented for the study. 29 were excluded and 104 patients were included with 57 of them receiving IV acetaminophen and 47 receiving placebo. There were no differences in baseline demographic characteristics including patient age, body mass index, gravidity, parity, ethnicity, comorbidities, or number of prior cesarean sections. There were no differences in estimated blood loss and length of surgery between groups. The total amount of narcotic medications consumed by patients receiving IV acetaminophen were significantly reduced when compared to the placebo group (47mg versus 65mg of oxycodone, p -value= 0.034). The total amount of ibuprofen used between groups was not different. There was no difference in the pain scores between the two groups before and after dose administration. There were no significant differences for reported medication side effects (nausea/emetis, respiratory depression, constipation) in either study arm.

CONCLUSION: IV acetaminophen in the post-operative time period for cesarean section surgery resulted in a significant decrease in the need for narcotic medications required for pain control.

34 Effect of a randomized controlled trial of an intensive medically supervised exercise program designed to improve maternal glucose control on gestational weight gain



Niamh Daly, Maria Farren, Aoife McKeating, Amy O’Higgins, Laura Mullaney, Michael J. Turner

UCD Centre for Human Reproduction, Coombe Women and Infants University Hospital, Dublin 8, Ireland

OBJECTIVE: To evaluate whether women with a body mass index (BMI) $>29.9\text{kg/m}^2$ enrolled in an intensive medically-supervised exercise intervention demonstrate a lower incidence of excessive gestational weight gain (GWG) compared to women undergoing routine care.

STUDY DESIGN: This is a prospective single-blinded randomized trial of obese women. Exclusion criteria were pre-existing diabetes mellitus, medications, no English, and BMI $<30\text{kg/m}^2$. Women were randomly assigned to: (1) routine prenatal care or (2) medically-supervised exercise intervention including three supervised exercise classes per week, invitation to a secret Facebook group to create a sense of community among participants and deliver healthy lifestyle advice. The study was adequately powered to detect a 7.2g/dL (0.4mmol/L) difference in mean fasting plasma glucose (FPG) levels on 2hOGTT. BMI and body composition were measured using bioelectrical impedance (Tanita MF 180CA) at the first prenatal visit and serially at 24-28 and 36 weeks gestation and at 6 weeks postpartum. Third trimester GWG was calculated at 36 weeks gestation. FPG was also measured. Excessive GWG was defined as $>9\text{kg}$ for obese women as recommended by the Institute of Medicine (IOM). Independent t-test and chi-squared were used where appropriate.

RESULTS: Of 88 women randomized, 44 were assigned to intervention and 44 received routine care. Maternal age, race, parity, initial BMI were similar between groups (all $P>0.05$). There was no difference in mean 3rd trimester FPG between groups ($p=0.66$). The follow up rates were similar (95.5% v 85.7%; $P=0.09$) between groups. Anthropometric measurements are shown in Table 1. Excessive gestational weight gain was lower in the intervention group (22.2% v 43.2%; $P<0.001$).

CONCLUSION: Women with a BMI $>29.9\text{kg/m}^2$ enrolled in this medically-supervised exercise intervention program demonstrate a lower incidence of excessive gestational weight gain at 36 weeks’ gestation. The same effect, however, is not seen on the mean GWG or mean fasting plasma glucose results. Clinicians should focus on improving pre-pregnancy BMI in this high risk population.

Table 1: Anthropometric changes during pregnancy and at 6 weeks postpartum

Anthropometrics	Early pregnancy		P-value	24-28 weeks gestation		P-value	36 weeks gestation		P-value	6 wks postpartum		P-value
	Control (n=46)	Intervention (n=42)		Control (n=43)	Intervention (n=39)		Control (n=42)	Intervention (n=39)		Control (n=36)	Intervention (n=31)	
Mean weight (kg)	95.8(15.8)	93.9(14.8)	0.56	101.1(15.4)	98.8(13.7)	0.49	103.6(15.6)	101.1(13.9)	0.46	96.7(16.2)	91.9(14.1)	0.09
Mean BMI (kg/m ²)	35.0(5.2)	34.7(4.7)	0.82	36.8(5.2)	36.5(4.5)	0.78	37.8(5.3)	37.4(4.7)	0.71	35.4(5.9)	34.0(4.7)	0.27
Mean Fat %	41.7(4.0)	41.8(4.0)	0.96	42.6(3.8)	42.4(3.3)	0.80	42.2(3.9)	41.6(3.4)	0.47	40.8(4.3)	40.2(4.0)	0.57
Mean visceral fat level	8.5(2.9)	8.5(2.4)	1.00	9.2(2.8)	9.1(2.1)	0.84	9.5(2.8)	9.2(2.1)	0.60	8.6(3.1)	8.1(2.3)	0.40
Mean gestational weight gain	n/a	n/a	n/a	4.9(3.8)	3.8(3.9)	0.19	7.9(4.8)	6.2(4.0)	0.13	5.2(5.4)	-1.6(1.2)	0.22
Excessive weight gain (IOM)							19 (43.2%)	8 (22.2%)	<0.001			