

(Dr Cox), University of Toronto, Toronto, Ontario, Canada; the Departments of Cellular and Molecular Medicine (Drs Benton and Bainbridge) and Pathology and Laboratory Medicine (Dr Gynspan), and the Interdisciplinary School of Health Sciences (Dr Bainbridge), University of Ottawa, Ottawa, Ontario, Canada.

<sup>1</sup>These authors contributed equally to this article.

Supported by the Canadian Institutes of Health Research Grant #128369 (S.A.B. and B.J.C.), an Undergraduate Summer Research Award from the University of Toronto (Department of Physiology; I.G.), an Ontario Graduate Scholarship (K.L.), and a Tier 2 Canada Research Chair in Placental Development and Maternal-Fetal Health (B.J.C.). S.J.B. is a Molly Towell Research Fellow.

The funding sources played no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

The authors report no conflict of interest.

Corresponding author: Brian J. Cox, PhD. [b.cox@utoronto.ca](mailto:b.cox@utoronto.ca)

---

## CORRECTION

**June 2018 (vol. 218, no. 6, pages 618.e1-7)**



Einerson BD, Rodriguez CE, Kennedy AM, et al. Magnetic resonance imaging is often misleading when used as an adjunct to ultrasound in the management of placenta accreta spectrum disorders. *Am J Obstet Gynecol* 2018;218:618.e1-7.

A Letter to the Editors in this issue identifies inconsistencies between the text and a figure in the reporting of ultrasound interpretation data in the article cited above. The authors of the article provide clarification.

The figure in question, Figure 1, is a flowchart that appears only in the online edition of the Journal, not in the paper edition.