

# Clinical diagnosis of endometriosis: a call to action



Sanjay K. Agarwal, MD; Charles Chapron, MD; Linda C. Giudice, MD, PhD; Marc R. Laufer, MD; Nicholas Leyland, MD; Stacey A. Missmer, ScD; Sukhbir S. Singh, MD; Hugh S. Taylor, MD

Endometriosis has such wide-ranging and pervasive sequelae that it has been described as “nothing short of a public health emergency” requiring immediate action.<sup>1</sup> Population-based data suggest that more than 4 million reproductive-age women have diagnosed endometriosis in the United States.<sup>2</sup> As daunting as this number is, it only tells part of the story, as an estimated 6 of 10 endometriosis cases are undiagnosed.<sup>3</sup> Thus more than 6 million American women may experience repercussions of endometriosis without the benefit of understanding the cause of their symptoms or appropriate management.

When discussing the patient’s experience with endometriosis, pain and infertility are usually of greatest concern, as they are 2 of the disease’s more common symptoms. However, the real toll is even greater: women with endometriosis experience diminished quality of life, increased incidence of depression, adverse effects on intimate relationships,

**THE PROBLEM:** Endometriosis is undiagnosed in a large proportion of affected women, resulting in ongoing and progressive symptoms with associated negative impacts on health and well-being. Current practice standards, which rely primarily on laparoscopy for a definitive diagnosis before beginning therapy, frequently result in prolonged delay between symptom onset, diagnosis, and subsequent treatment.

**A SOLUTION:** Enhanced use of clinical diagnostic techniques may reduce the delay in time to diagnosis and hence bring more rapid relief to affected patients, limit disease progression, and prevent sequelae.

limitations on participation in daily activities, reduced social activity, loss of productivity and associated income, increased risk of chronic disease, and significant direct and indirect healthcare costs.<sup>4–8</sup> Moreover, emerging data indicate that endometriosis is associated with greater risk of obstetric and neonatal complications.<sup>9–12</sup>

## The challenge of diagnosing endometriosis

There are no pathognomonic features or biomarkers necessary and sufficient to

define endometriosis. Rather, key symptoms that currently prompt surgical evaluation, such as pain and infertility, can have multiple causes. Endometriosis is typically defined by its histology: extrauterine lesions consisting of endometrial glands, endometrial stroma, and/or hemosiderin-laden macrophages. Based on location and depth, lesions are further described as superficial peritoneal lesions, ovarian endometrioma, or deep endometriosis. However, the presence of lesions does not preclude other etiologies for the

From the Center for Endometriosis Research and Treatment (Dr Agarwal), University of California San Diego, La Jolla, CA; Université Paris Descartes (Dr Chapron), Sorbonne Paris Cité, Faculté de Médecine, Assistance Publique—Hôpitaux de Paris (AP-HP), Hôpital Universitaire Paris Centre (HUPC), Centre Hospitalier Universitaire (CHU) Cochin, Department of Gynecology Obstetrics II and Reproductive Medicine, Paris, France; Department of Obstetrics, Gynecology and Reproductive Sciences (Dr Giudice), University of California, San Francisco, CA; Boston Center for Endometriosis, Boston Children’s Hospital and Brigham and Women’s Hospital; Department of Obstetrics, Gynecology, and Reproductive Biology (Dr Laufer), Brigham and Women’s Hospital and Harvard Medical School; Division of Gynecology, Boston Children’s Hospital and Harvard Medical School, Boston, MA; Department of Obstetrics and Gynaecology (Dr Leyland), McMaster University, Hamilton, Ontario, Canada; Boston Center for Endometriosis, Boston Children’s Hospital and Brigham and Women’s Hospital; Department of Epidemiology, Harvard T.H. Chan School of Public Health (Dr Missmer), Boston, MA; and Department of Obstetrics, Gynecology, and Reproductive Biology, Michigan State University, Grand Rapids, MI; Department of Obstetrics and Gynaecology (Dr Singh), University of Ottawa, and Ottawa Hospital Research Institute, Ottawa, Ontario, Canada; Department of Obstetrics, Gynecology, and Reproductive Sciences (Dr Taylor), Yale School of Medicine, New Haven, CT.

Received Aug. 24, 2018; revised Dec. 21, 2018; accepted Dec. 31, 2018.

S.K.A. is a consultant for and has received research support from AbbVie. C.C. is consultant for Ipsen, Bayer, AbbVie, and Gedeon Richter. He is an advisor/consultant for AbbVie, Bayer, Ipsen, and Gedeon Richter Pregel. L.C.G. is a consultant for AbbVie, Myovant Sciences, ForEndo Pharma, NextGen Jane, and Merck. M.R.L. is a consultant for AbbVie and NextGen Jane. N.L. is a consultant for and has received research support from AbbVie, Allergan, and the Canadian Institutes for Health Research. S.A.M. is a consultant for AbbVie, Oratel Diagnostics, and Celmatix, and receives research support from the National Institutes of Health and the Marriott Family Foundations. S.S.S. is a consultant for and has received research support from AbbVie, Bayer, and Allergan. H.S.T. is a consultant for AbbVie, Bayer, Obseva, OvaScience, ForEndo, and DotLab.

Support for the development of this manuscript was provided by AbbVie, Inc. AbbVie had the opportunity to review the final manuscript draft, but the manuscript content was solely at the discretion of the authors and reflects the opinions of the authors. The authors received no direct compensation for their efforts.

Corresponding author: Hugh S. Taylor, MD. [hugh.taylor@yale.edu](mailto:hugh.taylor@yale.edu)

0002-9378

© 2019 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).  
<https://doi.org/10.1016/j.ajog.2018.12.039>

patient's symptoms, and the lack of obvious lesions does not eliminate the possibility of endometriosis. Furthermore, there is poor correlation between symptoms and severity or extent of disease, as quantified by current staging systems.<sup>13</sup> From a clinical perspective, endometriosis may be better defined as a menstrual cycle–dependent, chronic, inflammatory, systemic disease that commonly presents as pelvic pain. Moving from a histological to a clinical definition opens the door to a different approach to diagnosis, one that emphasizes symptoms and their origins over lesion presence or absence, and that may, in the future, be validated by specific, noninvasive disease biomarkers.

Among those who ultimately receive a successful definitive diagnosis, contemporary literature describes delays from symptom onset to diagnosis ranging from 4 to 11 years.<sup>5,14–18</sup> Several factors exacerbate this delay,<sup>14,15,17,18</sup> including “normalization” of symptoms and misdiagnosis.<sup>15</sup> The presence of diagnostic delays is a worldwide phenomenon, occurring even in countries with universal healthcare.<sup>15,17</sup> Consequences of the delay in diagnosis are experienced by patients in multiple ways, including persistent symptoms and a commensurate detrimental impact on quality of life,<sup>14</sup> erosion of the patient–physician relationship,<sup>4,5</sup> and development of central sensitization—a mechanism whereby persistent endometriosis-associated pain increases pain awareness, even at sites unconnected anatomically with the lesion(s).<sup>14,19–21</sup> Moreover, although the evidence is limited, failure of timely diagnosis and adequate endometriosis management may foster disease progression and adhesion formation that may compromise fertility and increase the risk of central sensitization and chronic pelvic pain.<sup>22–24</sup>

The current diagnostic paradigm, endorsed by professional societies, requires laparoscopy with or without histologic verification as the gold standard, although many societies endorse the treatment of symptoms before obtaining a definitive surgical diagnosis.<sup>25–29</sup> Notably, the 2017 National Institute for

Health and Care Excellence guidelines reflect a philosophical shift, presenting empiric therapy prior to laparoscopy in the diagnostic and treatment algorithm unless fertility is a priority.<sup>30</sup> Although the merits of laparoscopy and its role in disease management should not be minimized, its accuracy, risks, and cost-effectiveness warrant reevaluation. The poor correlation between reported symptoms and extent of disease found at laparoscopy further illustrates the limitations of surgical disease assessment.<sup>31</sup> Detecting endometriosis via laparoscopy relies on the visual identification of lesions, a practice that is challenged by heterogeneous lesion appearance,<sup>32</sup> inaccessible lesion location (particularly for deep lesions),<sup>33</sup> and interobserver variability.<sup>34</sup> Surgical risks associated with laparoscopy are generally low,<sup>33,35</sup> although they merit consideration, given the potential for major (albeit rare) complications<sup>36</sup> and the need for re-treatment after initial laparoscopy because there is no surgical cure for endometriosis.<sup>37</sup> From a pragmatic perspective, evaluation of laparoscopy for endometriosis diagnosis and management must include a discussion of costs, which are substantially higher compared with nonsurgical approaches.<sup>38</sup>

### Argument for clinical diagnosis

Reliance on laparoscopy for endometriosis diagnosis supports the viewpoint that the presence of identifiable lesions in the pelvis is the central tenet of endometriosis, rather than approaching endometriosis as a menstrual cycle–dependent, chronic, inflammatory, systemic disease that often presents as pelvic pain. By shifting the paradigm to the patient rather than the lesion, the path to clinical diagnosis has the potential to be more inclusive with reduced diagnostic delay. Indeed, Soliman et al<sup>18</sup> reported diagnosing endometriosis by nonsurgical methods shortened the mean time from first consultation to diagnosis compared with surgical diagnosis. This shift, however, requires clinical diagnostic methodologies that accurately identify endometriosis. To that end, we have

compiled data on the accuracy of clinical assessments for diagnosing endometriosis (Table 1). Notably, these studies were highly heterogeneous, which precluded performance of a meaningful meta-analysis.

**Symptoms.** Pelvic pain, although common among women with endometriosis, is insufficient alone as an indicator of endometriosis, as it can be associated with several gynecologic (and non-gynecologic) conditions.<sup>39</sup> However, pelvic pain that is described as chronic, cyclic, and persistent or progressive (ie, worsening over time) increases the likelihood of an association with endometriosis.<sup>2,40,41</sup> Pain is typically initially menstrual (dysmenorrhea), but may progress to include nonmenstrual pelvic pain, which is prevalent among women with diagnosed endometriosis.<sup>42</sup> When asked about their experiences living with endometriosis, participants in the qualitative study by Moradi et al<sup>5</sup> universally described their pain as “severe and progressive during menstrual and nonmenstrual phases.” Women with endometriosis are more likely to report dyspareunia, dyschezia, and dysuria than unaffected women.<sup>2,40,43–46</sup> Although the sensitivity of dyspareunia is generally low,<sup>47–49</sup> indicating that its presence is not specific to endometriosis, deep dyspareunia is associated with deep endometriosis.<sup>46</sup>

Response of pain to treatment may be another indicator of endometriosis. Although nonsteroidal anti-inflammatory drugs (NSAIDs) effectively treat primary dysmenorrhea, pain reduction with these agents may be insufficient in women with endometriosis.<sup>26,28</sup> However, caution is indicated before dismissing NSAID-responsive pain as simply dysmenorrhea; early symptoms of endometriosis may be responsive to these agents, and we should not miss an opportunity to treat the disease before the development of serious sequelae.

**Patient and family history.** History of infertility is strongly associated with

TABLE 1

## Predictive value of signs, symptoms, and clinical findings for diagnosing endometriosis

Study design and population	Method of diagnosis	Assessment or parameter	Results
Endometriosis (general)			
Saha 2017 <sup>47a</sup> Cross-sectional survey of a Swedish twin cohort (N = 26,898)	Endometriosis diagnosis listed in electronic medical record	Severe dysmenorrhea	Sensitivity, 58%; specificity, 70%
		Chronic pelvic pain	Sensitivity, 25%; specificity, 89%
		Dyspareunia	Sensitivity, 16%; specificity, 96%
		Infertility	Sensitivity, 28%; specificity, 93%
		Oral pill as contraceptive	Sensitivity, 16%; specificity, 80%
Fuldeore 2017 <sup>2</sup> Respondents to an online, cross-sectional survey (N = 48,020)	Self-report (replying in the affirmative that a doctor had previously told the subject that she has or is suspected of having endometriosis)	Menstrual pelvic pain/cramping	OR, 1.6 (95% CI, 1.4–1.8)
		Nonmenstrual pelvic pain/cramping	OR, 4.1 (95% CI, 3.6–4.6)
		Dyspareunia	OR, 3.1 (95% CI, 2.8–3.5)
		Heavy menstrual bleeding	OR, 1.5 (95% CI, 1.3–1.7)
		Excessive or irregular bleeding	OR, 2.1 (95% CI, 1.8–2.4)
		Passage of clots	OR, 1.8 (95% CI, 1.6–2.0)
		Irregular menstrual periods (timing/duration)	OR, 1.5 (95% CI, 1.3–1.7)
		Constipation/bloating/diarrhea	OR, 1.9 (95% CI, 1.7–2.2)
		Fatigue/weariness/anemia	OR, 2.2 (95% CI, 2.0–2.5)
		Infertility	OR, 3.6 (95% CI, 3.0–4.4)
Ashrafi 2016 <sup>50</sup> Retrospective case-control study involving women who underwent laparoscopy for infertility evaluation (341 with endometriosis; 332 with a normal pelvis)	Laparoscopically visualized endometriosis	Family history of endometriosis	OR, 2.7 (95% CI, 1.06–7.1)
		History of galactorrhea	OR, 1.8 (95% CI, 1.1–3.05)
		History of pelvic surgery	OR, 14.5 (95% CI, 6.1–34.2)
		Dysmenorrhea	OR, 1.8 (95% CI, 1.1–2.8)
		Pelvic pain	OR, 4.1 (95% CI, 2.4–6.8)
		Dyspareunia	OR, 1.6 (95% CI, 1.09–2.4)
		Premenstrual spotting	OR, 2.2 (95% CI, 1.3–3.6)
		Fatigue	OR, 2.6 (95% CI, 1.3–5.1)
Apostolopoulos 2016 <sup>64</sup> Prospective, observational study of women who underwent laparoscopy for chronic pelvic pain (N = 144)	Laparoscopically visualized endometriosis	Noncyclical pain	Endometriosis, 62.5%; no endometriosis, 70.8%; <i>p</i> = 0.48
		Dysmenorrhea	Endometriosis, 79.1%; no endometriosis, 87.5%; <i>p</i> = 0.37
		Dyspareunia	Endometriosis, 25.0%; no endometriosis, 33.3%; <i>p</i> = 0.46
		Dyschezia	Endometriosis, 25.0%; no endometriosis, 20.8%; <i>p</i> = 0.69

Agarwal. Clinical diagnosis of endometriosis. *Am J Obstet Gynecol* 2019.

(continued)

endometriosis, although this may be skewed due to more thorough evaluation of women with infertility increasing the chances of successful diagnosis.<sup>2,41,43-47</sup>

Other factors associated with a greater likelihood of successful endometriosis diagnosis are family history of the disease,<sup>43,50</sup> previous pelvic surgery,<sup>50</sup>

and a history of benign ovarian cysts and/or ovarian pain.<sup>43,45</sup>

**Menstrual cycle characteristics.** In a recent cross-sectional survey of approximately 50,000 women, several menstrual cycle characteristics were more prevalent among women with vs

without diagnosed endometriosis, including heavy menstrual bleeding, excessive/irregular bleeding, passing clots, and irregular menstrual periods. Premenstrual spotting also correlates with endometriosis in infertile women.<sup>48,50</sup> Although these disorders are common in women with

**TABLE 1**  
**Predictive value of signs, symptoms, and clinical findings for diagnosing endometriosis** (continued)

Study design and population	Method of diagnosis	Assessment or parameter	Results
Schliep 2015 <sup>40</sup> Operative cohort from the ENDO study—women without a history of surgically confirmed endometriosis who underwent laparoscopy or laparotomy (N = 473)	Surgically visualized endometriosis	Chronic pelvic pain	Endometriosis, 44.2%; other, 39.0%; normal pelvis, 30.2%; $p = 0.04$
		Cyclic pelvic pain	Endometriosis, 49.5%; other, 31.0%; normal pelvis, 33.1%; $p < 0.001$
		Vaginal pain with intercourse	Endometriosis, 54.7%; other, 41.5%; normal pelvis, 32.4%; $p < 0.001$
		Deep pain with intercourse	Endometriosis, 53.2%; other, 38.1%; normal pelvis, 30.9%; $p < 0.001$
		Burning vaginal pain after intercourse	Endometriosis, 33.2%; other, 22.5%; normal pelvis, 22.1%; $p = 0.03$
		Pain just before menstrual period	Endometriosis, 75.3%; other, 61.9%; normal pelvis, 66.2%; $p = 0.03$
		Level of cramps with period	Endometriosis, 91.1%; other, 85.0%; normal pelvis, 79.4%; $p = 0.01$
		Pain after period is over	Endometriosis, 38.4%; other, 26.5%; normal pelvis, 38.2%; $p = 0.04$
		Pain at ovulation (mid-cycle)	Endometriosis, 67.4%; other, 49.0%; normal pelvis, 52.2%; $p = 0.001$
		Dysuria	Endometriosis, 22.6%; other, 19.1%; normal pelvis, 11.0%; $p = 0.03$
Heitman 2014 <sup>48</sup> Retrospective cohort of consecutive women with or without pelvic pain who were evaluated for infertility (N = 80)	Histologically verified endometriosis	Premenstrual spotting for $\geq 2$ days	Sensitivity, 76%; specificity, 90%; PPV, 96%; NPV, 74%; accuracy, 81%
		Dysmenorrhea	Sensitivity, 87%; specificity, 63%; PPV, 75%; NPV, 79%; accuracy, 76%
		Dyspareunia	Sensitivity, 38%; specificity, 83%; PPV, 74%; NPV, 51%; accuracy, 58%

Agarwal. Clinical diagnosis of endometriosis. Am J Obstet Gynecol 2019.

(continued)

endometriosis, most of these women have regular cycles without abnormal bleeding.

**Physical examination.** Data from comparative studies suggest that findings on physical examination can identify

endometriosis with high accuracy.<sup>51-53</sup> For example, using defined criteria for a positive bimanual pelvic examination (ie, palpable nodularity, stiffened and/or thickened pelvic anatomy, especially the uterosacral ligaments, vagina, rectovaginal space, pouch of Douglas,

adnexa, rectosigmoid, or posterior wall of the urinary bladder), Hudelist et al<sup>51</sup> reported endometriosis diagnosis accuracy of 86–99%, depending on anatomic location. Diagnostic acumen of pelvic examination is lower for deep endometriosis,<sup>52,53</sup> although

**TABLE 1**  
**Predictive value of signs, symptoms, and clinical findings for diagnosing endometriosis** (continued)

Study design and population	Method of diagnosis	Assessment or parameter	Results
Peterson 2013 <sup>41b</sup> ENDO Study—Prospective, matched-exposure cohort study comprising women undergoing pelvic surgery (n = 495) and a matched cohort (n = 131)	Surgically visualized endometriosis (operative cohort) Pelvic MRI-diagnosed endometriosis (matched cohort)	History of infertility	OR, 2.43 (95% CI, 1.57–3.76) [operative]; 7.91 (1.69–37.2) [matched]
		Dysmenorrhea	OR, 2.46 (95% CI, 1.28–4.72) [operative]; 1.41 (0.28–7.14) [matched]
		Pelvic pain	OR, 1.39 (95% CI, 0.95–2.04) [operative]; 0.76 (0.09–6.54) [matched]
		Pelvic pain (surgical indication)	OR, 3.67 (95% CI, 2.44–5.50) [operative]
Nnoaham 2012 <sup>43</sup> Prospective, observational study of symptomatic women with scheduled laparoscopy (N = 1396)	Laparoscopically visualized endometriosis	Model comprising multiple factors (eg, dysmenorrhea, dyschezia, nonmenstrual pelvic pain, ovarian cyst, family history, race, etc)	Sensitivity, 85%; specificity, 44%
		Model and ultrasound	Sensitivity, 58%; specificity, 89%
Paulson 2011 <sup>54</sup> Prospective cohort of women with chronic pelvic pain (N = 284)	Laparoscopically or histologically confirmed endometriosis	Anterior vaginal wall tenderness (endometriosis and other pathology)	Sensitivity, 93%
		Anterior vaginal wall tenderness (endometriosis only)	Sensitivity, 17%
Droz 2011 <sup>65</sup> Retrospective cohort of women evaluated for chronic pelvic pain (N = 331)	Histologically verified endometriosis	Short-form MPQ pain descriptor:	
		Cramping	Sensitivity, 92%; specificity, 33%; PPV, 40%; NPV, 89%
		Sickening	Sensitivity, 73%; specificity, 46%; PPV, 40%; NPV, 78%
		Tiring/exhausting	Sensitivity, 77%; specificity, 38%; PPV, 38%; NPV, 77%
		Shooting	Sensitivity, 70%; specificity, 43%; PPV, 37%; NPV, 75%
		Punishing/cruel	Sensitivity, 49%; specificity, 65%; PPV, 40%; NPV, 72%
Paulson 2009 <sup>55</sup> Prospective study of consecutive women with unexplained infertility (N = 55)	Laparoscopically or histologically confirmed endometriosis	Anterior vaginal wall tenderness	Sensitivity, 84%; specificity, 75%; PPV, 86%; NPV, 69%
Meuleman 2009 <sup>39</sup> Retrospective case series comprising infertile women with regular cycles and no prior endometriosis diagnosis (N = 221)	Histologically verified endometriosis	Pelvic pain	Sensitivity, 59%; specificity, 56%; PPV, 54%; NPV, 57%
		Pelvic pain and type of infertility, age, and duration of infertility	Sensitivity, 65%; specificity, 73%
Hudelist 2009 <sup>51c</sup> Prospective study of consecutive women with symptoms of endometriosis (N = 200)	Histologically verified endometriosis	Vaginal examination	Sensitivity, 23–88%; specificity, 89–100%; PPV, 65–100%; NPV, 85–99%; accuracy, 86–99%
		Vaginal examination and TVS	Sensitivity, 67–100%; specificity, 86–100%; PPV, 50–100%; NPV, 93–100%; accuracy, 86–100%

**TABLE 1**  
**Predictive value of signs, symptoms, and clinical findings for diagnosing endometriosis** (continued)

Study design and population	Method of diagnosis	Assessment or parameter	Results
Flores 2008 <sup>44</sup> Respondents to a self-administered questionnaire (N = 1285)	Self-reported surgically confirmed endometriosis	Dysmenorrhea	Cases, 82.5%; general population, 59.3%; $p < 0.001$
		Severe dysmenorrhea	Cases, 65.9%; general population, 52.9%; $p = NS$
		Dyspareunia	Cases, 52.0%; general population, 20.0%; $p < 0.001$
		Problems conceiving	Cases, 70.6%; general population, 25.2%; $p < 0.001$
		Chronic pelvic pain	Cases, 80.0%; general population, 22.9%; $p < 0.001$
Ballard 2008 <sup>45d</sup> National case-control study comprising women with endometriosis (n = 5540) and matched controls (n = 21,239)	Diagnostic or procedural codes consistent with endometriosis recorded in a nationwide general practice database	Dysmenorrhea	OR, 9.8 (95% CI, 8.8–10.9)
		Pelvic pain	OR, 13.5 (95% CI, 11.7–15.7)
		Dyspareunia	OR, 9.4 (95% CI, 8.0–11.1)
		Abdominal pain	OR, 5.9 (95% CI, 5.5–6.4)
		Menorrhagia	OR, 5.0 (95% CI, 4.6–5.5)
		Intermenstrual pain	OR, 6.9 (95% CI, 4.7–10.2)
		Infertility/subfertility	OR, 6.2 (95% CI, 5.4–7.1)
		Pelvic inflammatory disease	OR, 6.4 (95% CI, 5.6–7.4)
		Ovarian cysts	OR, 12.2 (95% CI, 9.9–15.0)
		Ovary pain	OR, 9.1 (95% CI, 3.2–26.0)
Endometriosis (stages III and IV)			
Peterson 2013 <sup>41b</sup> ENDO Study—prospective, matched exposure cohort study comprising women undergoing pelvic surgery (n = 495)	Surgically visualized endometriosis (operative cohort)	History of infertility	OR, 4.74 (95% CI, 2.57–8.75)
		Dysmenorrhea	OR, 3.43 (95% CI, 1.02–11.5)
		Pelvic pain	OR, 1.60 (95% CI, 0.89–2.87)
		Pelvic pain (surgical indication)	OR, 4.47 (95% CI, 2.39–8.38)
Nnoaham 2012 <sup>43</sup> Prospective, observational study of symptomatic women with scheduled laparoscopy (N = 1396)	Laparoscopically visualized endometriosis	Model comprising multiple factors (eg, dyschezia, ovarian cyst, infertility, cycle length, GI/bladder symptoms, race, etc)	Sensitivity, 71%; specificity, 85%
		Model with ultrasound	Sensitivity, 82%; specificity, 76%
Endometriosis and other peri-ovarian adhesions			

Agarwal. Clinical diagnosis of endometriosis. *Am J Obstet Gynecol* 2019.

(continued)

examination during menses improves detection.<sup>26</sup> Anterior vaginal wall tenderness has low sensitivity for detecting endometriosis in women with chronic pelvic pain,<sup>54</sup> but demonstrates prognostic value for endometriosis among women with unexplained infertility.<sup>55</sup> A caveat to bimanual examination is that it may not be feasible for non–sexually active adolescents/young adults and may not identify early-stage, superficial disease.

**Combination assessments.** The ability to identify endometriosis nonsurgically is enhanced when multiple factors are combined. Ballard et al<sup>45</sup> reported that the likelihood of endometriosis increased with the number of symptoms present, from an odds ratio of 5.0 with 1 symptom to 84.7 for 7 or more symptoms. Several investigators have used this approach to develop models for predicting endometriosis.<sup>43,46,56</sup> Using data from a

prospective, multinational study, Nnoaham et al<sup>43</sup> created a model combining symptoms and patient history with ultrasound findings that predicted revised American Society for Reproductive Medicine (rASRM) stage III and IV endometriosis with good accuracy. The authors suggest that such screening tools could reduce “diagnostic delay, high investigation costs, and personal suffering associated with endometriosis.”<sup>43</sup>

**TABLE 1**  
**Predictive value of signs, symptoms, and clinical findings for diagnosing endometriosis** (continued)

Study design and population	Method of diagnosis	Assessment or parameter	Results
Marasinghe 2014 <sup>49</sup> Prospective, observational study comprising women evaluated for infertility and/or chronic pelvic pain (N = 110)	Laparoscopically visualized endometriosis	Dyspareunia	Sensitivity, 46%; specificity, 77%; PPV, 52%; NPV, 73%; accuracy, 47%
		Dysmenorrhea	Sensitivity, 76%; specificity, 70%; PPV, 57%; NPV, 84%; accuracy, 71%
		Dyspareunia and dysmenorrhea	Sensitivity, 78%; specificity, 64%; PPV, 54%; NPV, 85%; accuracy, 68%
		Vaginal examination	Sensitivity, 73%; specificity, 88%; PPV, 77%; NPV, 86%; accuracy, 83%
		Dyspareunia, dysmenorrhea, and vaginal examination	Sensitivity, 84%; specificity, 62%; PPV, 54%; NPV, 88%; accuracy, 69%
		Fixed ovaries on TVS	Sensitivity, 78%; specificity, 94%; PPV, 88%; NPV, 89%; accuracy, 88%
Dyspareunia, dysmenorrhea, vaginal examination and fixed ovaries			Sensitivity, 92%; specificity, 61%; PPV, 56%; NPV, 93%; accuracy, 71%
<b>Deep endometriosis</b>			
Perello 2017 <sup>56</sup> Retrospective analysis of consecutive women with ovarian endometrioma who underwent surgery (N = 178)	Histologically verified endometriosis	Model including previous pregnancy, history of surgery for endometriosis, endometriosis-associated pelvic pain score	Sensitivity, 80%; specificity, 84%
Lafay Pillet 2014 <sup>46e</sup> Prospective, single-center study of women with a histological diagnosis of endometriosis (N = 211)	Histologically verified endometriosis	Infertility (primary or secondary)	Sensitivity, 51%; specificity, 73%; OR, 1.5; <i>p</i> = 0.003
		Duration of pain >24 mo	Sensitivity, 62%; specificity, 81%; OR, 7.1; <i>p</i> < 0.001
		VAS deep dyspareunia >5	Sensitivity, 69%; specificity, 59%; OR, 3.2; <i>p</i> = 0.007
		VAS GI symptoms ≥5	Sensitivity, 75%; specificity, 76%; OR, 9.3; <i>p</i> < 0.001
		Severe dysmenorrhea	Sensitivity, 55%; specificity, 75%; OR, 3.5; <i>p</i> < 0.001
Hudelist 2011 <sup>52c</sup> Prospective study of premenopausal women with suspected endometriosis (N = 129)	Histologically verified endometriosis	Vaginal examination	Sensitivity, 25–78%; specificity, 80–100%; PPV, 43–100%; NPV, 84–98%; accuracy, 73–98%
		TVS	Sensitivity, 50–96%; specificity, 96–100%; PPV, 50–100%; NPV, 90–99%; accuracy, 90–99%

Agarwal. Clinical diagnosis of endometriosis. *Am J Obstet Gynecol* 2019.

(continued)

**Additional considerations.** Imaging can be a useful adjunct to clinical diagnostic measures, and transvaginal ultrasound improves accuracy when used adjunctively with symptoms, patient

history, and/or physical findings.<sup>43,49,51</sup> Ultrasound is particularly sensitive for detecting ovarian endometriomas and deep endometriosis.<sup>25,57,58</sup> Indeed, a Cochrane meta-analysis found that

transvaginal ultrasound approaches the sensitivity and specificity needed to replace surgery for endometrioma detection.<sup>57</sup> The International Deep Endometriosis Analysis (IDEA) group

**TABLE 1**  
**Predictive value of signs, symptoms, and clinical findings for diagnosing endometriosis** (continued)

Study design and population	Method of diagnosis	Assessment or parameter	Results
Bazot 2009 <sup>53c</sup> Retrospective, longitudinal study of consecutive women with clinical evidence of endometriosis (N = 92)	Laparoscopically visualized endometriosis	Vaginal examination	Sensitivity, 18–74%; specificity, 72–96%; PPV, 40–97%; NPV, 24–90%; accuracy, 54–87%
		TVS	Sensitivity, 9–94%; specificity, 67–100%; PPV, 50–100%; NPV, 25–89%; accuracy, 77–96%
		Rectal endoscopic sonography	Sensitivity, 7–89%; specificity, 44–100%; PPV, 33–100%; NPV, 9–90%; accuracy, 48–90%
		MRI	Sensitivity, 55–87%; specificity, 86–99%; PPV, 73–99%; NPV, 38–94%; accuracy, 84–94%


To identify relevant studies, a search of the MEDLINE database was performed using the following search terms: endometriosis AND (pain OR cycle OR infertility OR “physical exam” OR “physical examination” OR “pelvic exam” OR “pelvic examination”) AND (specificity OR sensitivity OR accuracy). Articles were limited to clinical studies published in English from 2008 through March 2018. Additional studies identified via citations in associated manuscripts were added if applicable.

ENDO, Endometriosis: Natural History, Diagnosis, and Outcomes Study; GI, gastrointestinal; HR, hazard ratio; MPQ, McGill Pain Questionnaire; MRI, magnetic resonance imaging; NPV, negative predictive value; PPV, positive predictive value; TVS, transvaginal sonography; VAS, visual analogue scale.

<sup>a</sup> Reported are the agreement between self-reported symptoms of endometriosis and diagnosis of endometriosis recorded in medical records; <sup>b</sup> Data are adjusted odds ratios; <sup>c</sup> Ranges reflect different values based on anatomic locations of the endometriotic lesions; <sup>d</sup> Shown here are symptoms and signs with an odds ratio for predicting endometriosis of 5.0 or greater; <sup>e</sup> Lafay Pillet et al<sup>18</sup> evaluated combining multiple signs, symptoms, and findings to predict the presence of deep endometriosis. Presented here are the individual measures included in the final model.

Agarwal. Clinical diagnosis of endometriosis. Am J Obstet Gynecol 2019.

**FIGURE 1**  
**Algorithm for a clinical diagnosis of endometriosis**



\*Alternative diagnoses indicated by symptoms on the right side of the chart may coexist with endometriosis and do not rule out the presence of endometriosis.

Agarwal. Clinical diagnosis of endometriosis. Am J Obstet Gynecol 2019.

consensus statement on systematic sonographic evaluation of the pelvis in women with suspected endometriosis provides standards for improved imaging.<sup>59</sup> Traditional routine transvaginal ultrasound may be limited to endometrioma diagnosis; however, “expert-guided” imaging, as outlined by the IDEA group, will help improve clinical assessment across endometriosis manifestations. Nonetheless, not all endometriosis will be visualized by imaging, and imaging cannot be used to rule out endometriosis.

Magnetic resonance imaging is a noninvasive option; however, it is expensive, not universally available, and lacks sensitivity, and is therefore infrequently used for endometriosis diagnosis. Although many are currently being studied, as yet, no noninvasive or minimally invasive biomarker has been established to diagnose endometriosis.<sup>60-62</sup>

Much of what is known about endometriosis comes from surgically diagnosed adults. Increased research into endometriosis among surgically diagnosed adolescents and prospective

studies of those with suggestive signs and symptoms will help to better identify hallmarks of disease onset and risk factors for disease progression and treatment prognosis. Although a detailed review of endometriosis in adolescents is beyond the scope of this discussion, it is noteworthy that endometriosis occurs in adolescents and that patients who are younger at the time of symptom onset experience longer diagnostic delays than older patients.<sup>17,18</sup> This delay is attributed to prolonged time before seeking treatment and a longer interval between first clinical consultation and referral or diagnosis. It is important that clinicians evaluate symptoms that merit suspicion in adolescents as seriously as in adults.<sup>42</sup>

### Implementing clinical diagnosis

Clinical diagnosis is already applied in clinical practice, albeit inconsistently and without standardization.<sup>2,18</sup> In an effort to provide a unified, practical approach to clinically diagnosing endometriosis, we have developed an algorithm informed by evidence in the literature and clinical experience (Figure 1). The proposed algorithm uses techniques readily available to most practitioners and allows clinicians to initiate treatment without delay or invasive procedures. For each step, we identify findings that are consistent with endometriosis and those suggesting a possible alternative diagnosis. In general, persistent and/or worsening cyclic or constant pelvic pain, particularly in the presence of other endometriosis-associated symptoms, patient history, and findings on physical examination, suggest endometriosis. When these findings are unclear, imaging with transvaginal ultrasound is a widely available and low-cost option.

This algorithm does not diminish the value of laparoscopy as a treatment option in those for whom medical therapy is insufficient, nor does it minimize laparoscopy as a diagnostic tool when clinical signs are uncertain or suggest nonendometriosis pathology (eg, other benign or malignant ovarian neoplasms). Rather, the algorithm is intended to make the diagnosis of endometriosis more accessible, reducing

the negative impact of undiagnosed and untreated endometriosis on women's lives. Practitioners should feel empowered to clinically diagnose this disease early and without an invasive procedure. Although the ramifications of early diagnosis and treatment have not been studied, the potential exists to relieve pain, to avoid central sensitization and pain persistence, to prevent infertility, and to change the trajectory of patients' lives. It is increasingly recognized that chronic diseases such as endometriosis generate cumulative life-course impairment through limitations imposed on life choices, including education, career, and family.<sup>5,63</sup> Overall patient health may also be improved by addressing the psychosocial and physical manifestations often found in conjunction with endometriosis, such as persistent pelvic pain, depression, anxiety, fatigue, bloating/weight gain, gastrointestinal issues, and sexual dysfunction.<sup>2,4,5,50</sup> Now is the time to change the paradigm of the diagnosis of endometriosis by increasing speed and validity, leading to improved access to effective early treatment. ■

### ACKNOWLEDGMENTS

Medical writing support for development of this manuscript, funded by AbbVie, Inc., was provided by Crystal Murcia, PhD, and Lamara D. Shrode, PhD, CMPP, of JB Ashtin.

### REFERENCES

- Hatch O. This is nothing short of a public health emergency. CNN. March 28, 2018. Available at: <https://www.cnn.com/2018/03/27/opinions/endometriosis-start-a-conversation-hatch-opinion/index.html>. Accessed April 14, 2018.
- Fuldeore MJ, Soliman AM. Prevalence and symptomatic burden of diagnosed endometriosis in the United States: national estimates from a cross-sectional survey of 59,411 women. *Gynecol Obstet Invest* 2017;82:453–61.
- Morassutto C, Monasta L, Ricci G, Barbone F, Ronfani L. Incidence and estimated prevalence of endometriosis and adenomyosis in northeast Italy: a data linkage study. *PLoS One* 2016;11:e0154227.
- Culley L, Law C, Hudson N, et al. The social and psychological impact of endometriosis on women's lives: a critical narrative review. *Hum Reprod Update* 2013;19:625–39.
- Moradi M, Parker M, Sneddon A, Lopez V, Ellwood D. Impact of endometriosis on women's

lives: a qualitative study. *BMC Womens Health* 2014;14:123.

- Kvaskoff M, Mu F, Terry KL, et al. Endometriosis: a high-risk population for major chronic diseases? *Hum Reprod Update* 2015;21:500–16.
- Soliman AM, Coyne KS, Gries KS, Castelli-Haley J, Snabes MC, Surrey ES. The effect of endometriosis symptoms on absenteeism and presenteeism in the workplace and at home. *J Manag Care Spec Pharm* 2017;23:745–54.
- Soliman AM, Surrey E, Bonafede M, Nelson JK, Castelli-Haley J. Real-world evaluation of direct and indirect economic burden among endometriosis patients in the United States. *Adv Ther* 2018;35:408–23.
- Harada T, Taniguchi F, Onishi K, et al. Obstetrical complications in women with endometriosis: a cohort study in Japan. *PLoS One* 2016;11:e0168476.
- Berlac JF, Hartwell D, Skovlund CW, Langhoff-Roos J, Lidgaard O. Endometriosis increases the risk of obstetrical and neonatal complications. *Acta Obstet Gynecol Scand* 2017;96:751–60.
- Zullo F, Spagnolo E, Saccone G, et al. Endometriosis and obstetrics complications: a systematic review and meta-analysis. *Fertil Steril* 2017;108:667–72.
- Chen I, Lalani S, Xie RH, Shen M, Singh SS, Wen SW. Association between surgically diagnosed endometriosis and adverse pregnancy outcomes. *Fertil Steril* 2018;109:142–7.
- Johnson NP, Hummelshoj L, Adamson GD, et al. World Endometriosis Society consensus on the classification of endometriosis. *Hum Reprod* 2017;32:315–24.
- Nnoaham KE, Hummelshoj L, Webster P, et al. Impact of endometriosis on quality of life and work productivity: a multicenter study across ten countries. *Fertil Steril* 2011;96:366–73.
- Hudelist G, Fritzer N, Thomas A, et al. Diagnostic delay for endometriosis in Austria and Germany: causes and possible consequences. *Hum Reprod* 2012;27:3412–6.
- Fourquet J, Sinaii N, Stratton P, et al. Characteristics of women with endometriosis from the USA and Puerto Rico. *J Endometr Pelvic Pain Disord* 2015;7:129–35.
- Staal AH, Van Der Zanden M, Nap AW. Diagnostic delay of endometriosis in the Netherlands. *Gynecol Obstet Invest* 2016;81:321–4.
- Soliman AM, Fuldeore M, Snabes MC. Factors associated with time to endometriosis diagnosis in the United States. *J Womens Health (Larchmt)* 2017;26:788–97.
- Issa B, Onon TS, Agrawal A, et al. Visceral hypersensitivity in endometriosis: a new target for treatment? *Gut* 2012;61:367–72.
- As-Sanie S, Harris RE, Harte SE, Tu FF, Neshewat G, Clauw DJ. Increased pressure pain sensitivity in women with chronic pelvic pain. *Obstet Gynecol* 2013;122:1047–55.
- Li T, Mamillapalli R, Ding S, et al. Endometriosis alters brain electro-physiology, gene

- expression and increased pain sensitization, anxiety, and depression in female mice. *Biol Reprod* 2018;99:349–59.
- 22.** Unger CA, Laufer MR. Progression of endometriosis in non-medically managed adolescents: a case series. *J Pediatr Adolesc Gynecol* 2011;24:e21–3.
- 23.** Brosens I, Gordts S, Benagiano G. Endometriosis in adolescents is a hidden, progressive and severe disease that deserves attention, not just compassion. *Hum Reprod* 2013;28:2026–31.
- 24.** Coxon L, Horne AW, Vincent K. Pathophysiology of endometriosis-associated pain: a review of pelvic and central nervous system mechanisms. *Best Pract Res Clin Obstet Gynaecol* 2018;51:53–67.
- 25.** Practice bulletin no. 114: management of endometriosis. *Obstet Gynecol* 2010;116:223–36.
- 26.** Leyland N, Casper R, Laberge P, Singh SS, Society of Obstetricians and Gynaecologists of Canada. Endometriosis: diagnosis and management. *J Obstet Gynaecol Can* 2010;32:S1–32.
- 27.** Johnson NP, Hummelshoj L, World Endometriosis Society Montpellier Consortium. Consensus on current management of endometriosis. *Hum Reprod* 2013;28:1552–68.
- 28.** Practice Committee of the American Society for Reproductive Medicine. Treatment of pelvic pain associated with endometriosis: a committee opinion. *Fertil Steril* 2014;101:927–35.
- 29.** Dunselman GA, Vermeulen N, Becker C, et al. ESHRE guideline: management of women with endometriosis. *Hum Reprod* 2014;29:400–12.
- 30.** National Institute for Health and Care Excellence. Endometriosis: diagnosis and management (NG73). London, United Kingdom; 2017. Available at: <http://nice.org.uk/guidance/ng73>. Accessed May 18, 2018.
- 31.** Vercellini P, Fedele L, Aimi G, Pietropaolo G, Consonni D, Crosignani PG. Association between endometriosis stage, lesion type, patient characteristics and severity of pelvic pain symptoms: a multivariate analysis of over 1000 patients. *Hum Reprod* 2007;22:266–71.
- 32.** Albee RB Jr, Sinervo K, Fisher DT. Laparoscopic excision of lesions suggestive of endometriosis or otherwise atypical in appearance: relationship between visual findings and final histologic diagnosis. *J Minim Invasive Gynecol* 2008;15:32–7.
- 33.** Singh SS, Suen MW. Surgery for endometriosis: beyond medical therapies. *Fertil Steril* 2017;107:549–54.
- 34.** Schliep KC, Chen Z, Stanford JB, et al. Endometriosis diagnosis and staging by operating surgeon and expert review using multiple diagnostic tools: an inter-rater agreement study. *Br J Obstet Gynecol* 2017;124:220–9.
- 35.** Surrey ES, Soliman AM, Yang H, Du EX, Su B. Treatment patterns, complications, and health care utilization among endometriosis patients undergoing a laparoscopy or a hysterectomy: a retrospective claims analysis. *Adv Ther* 2017;34:2436–51.
- 36.** Chapron C, Fauconnier A, Goffinet F, Breart G, Dubuisson JB. Laparoscopic surgery is not inherently dangerous for patients presenting with benign gynaecologic pathology. Results of a meta-analysis. *Hum Reprod* 2002;17:1334–42.
- 37.** Soliman AM, Du EX, Yang H, Wu EQ, Haley JC. Retreatment rates among endometriosis patients undergoing hysterectomy or laparoscopy. *J Womens Health (Larchmt)* 2017;26:644–54.
- 38.** Soliman AM, Taylor HS, Bonafede M, Nelson JK, Castelli-Haley J. Incremental direct and indirect cost burden attributed to endometriosis surgeries in the United States. *Fertil Steril* 2017;107:1181–90.
- 39.** Meuleman C, Vandenabeele B, Fieuws S, Spiessens C, Timmerman D, D'hooghe T. High prevalence of endometriosis in infertile women with normal ovulation and normospermic partners. *Fertil Steril* 2009;92:68–74.
- 40.** Schliep KC, Mumford SL, Peterson CM, et al. Pain typology and incident endometriosis. *Hum Reprod* 2015;30:2427–38.
- 41.** Peterson CM, Johnstone EB, Hammoud AO, et al. Risk factors associated with endometriosis: importance of study population for characterizing disease in the ENDO Study. *Am J Obstet Gynecol* 2013;208:451.
- 42.** Divasta AD, Vitonis AF, Laufer MR, Missmer SA. Spectrum of symptoms in women diagnosed with endometriosis during adolescence vs adulthood. *Am J Obstet Gynecol* 2018;218:324.
- 43.** Nnoaham KE, Hummelshoj L, Kennedy SH, Jenkinson C, Zondervan KT, World Endometriosis Research Foundation Women's Health Symptom Survey Consortium. Developing symptom-based predictive models of endometriosis as a clinical screening tool: results from a multicenter study. *Fertil Steril* 2012;98:692–701.
- 44.** Flores I, Abreu S, Abac S, Fourquet J, Laboy J, Rios-Bedoya C. Self-reported prevalence of endometriosis and its symptoms among Puerto Rican women. *Int J Gynaecol Obstet* 2008;100:257–61.
- 45.** Ballard KD, Seaman HE, De Vries CS, Wright JT. Can symptomatology help in the diagnosis of endometriosis? Findings from a national case-control study—part 1. *Br J Obstet Gynecol* 2008;115:1382–91.
- 46.** Lafay Pillet MC, Huchon C, Santulli P, Borghese B, Chapron C, Fauconnier A. A clinical score can predict associated deep infiltrating endometriosis before surgery for an endometrioma. *Hum Reprod* 2014;29:1666–76.
- 47.** Saha R, Marions L, Tornvall P. Validity of self-reported endometriosis and endometriosis-related questions in a Swedish female twin cohort. *Fertil Steril* 2017;107:174–8.
- 48.** Heitmann RJ, Langan KL, Huang RR, Chow GE, Burney RO. Premenstrual spotting of  $\geq 2$  days is strongly associated with histologically confirmed endometriosis in women with infertility. *Am J Obstet Gynecol* 2014;211:358.
- 49.** Marasinghe JP, Senanayake H, Saravanabhava N, Arambepola C, Condous G, Greenwood P. History, pelvic examination findings and mobility of ovaries as a sonographic marker to detect pelvic adhesions with fixed ovaries. *J Obstet Gynaecol Res* 2014;40:785–90.
- 50.** Ashrafi M, Sadatmahalleh SJ, Akhoond MR, Talebi M. Evaluation of risk factors associated with endometriosis in infertile women. *Int J Fertil Steril* 2016;10:11–21.
- 51.** Hudelist G, Oberwinkler KH, Singer CF, et al. Combination of transvaginal sonography and clinical examination for preoperative diagnosis of pelvic endometriosis. *Hum Reprod* 2009;24:1018–24.
- 52.** Hudelist G, Ballard K, English J, et al. Transvaginal sonography vs. clinical examination in the preoperative diagnosis of deep infiltrating endometriosis. *Ultrasound Obstet Gynecol* 2011;37:480–7.
- 53.** Bazot M, Lafont C, Rouzier R, Roseau G, Thomassin-Naggara I, Darai E. Diagnostic accuracy of physical examination, transvaginal sonography, rectal endoscopic sonography, and magnetic resonance imaging to diagnose deep infiltrating endometriosis. *Fertil Steril* 2009;92:1825–33.
- 54.** Paulson JD, Paulson JN. Anterior vaginal wall tenderness (AVWT) as a physical symptom in chronic pelvic pain. *JLS* 2011;15:6–9.
- 55.** Paulson JD. Correlation of anterior vaginal wall pain with endometriosis in infertile patients. *J Reprod Med* 2009;54:145–50.
- 56.** Perello M, Martinez-Zamora MA, Torres X, et al. Markers of deep infiltrating endometriosis in patients with ovarian endometrioma: a predictive model. *Eur J Obstet Gynecol Reprod Biol* 2017;209:55–60.
- 57.** Nisenblat V, Bossuyt PM, Farquhar C, Johnson N, Hull ML. Imaging modalities for the non-invasive diagnosis of endometriosis. *Cochrane Database Syst Rev* 2016;2:CD009591.
- 58.** Turocy JM, Benacerraf BR. Transvaginal sonography in the diagnosis of deep infiltrating endometriosis: a review. *J Clin Ultrasound* 2017;45:313–8.
- 59.** Guerriero S, Condous G, Van Den Bosch T, et al. Systematic approach to sonographic evaluation of the pelvis in women with suspected endometriosis, including terms, definitions and measurements: a consensus opinion from the International Deep Endometriosis Analysis (IDEA) group. *Ultrasound Obstet Gynecol* 2016;48:318–32.
- 60.** Cho S, Mutlu L, Grechukhina O, Taylor HS. Circulating microRNAs as potential biomarkers for endometriosis. *Fertil Steril* 2015;103:1252–60.
- 61.** Fassbender A, Burney RO, O DF, D'hooghe T, Giudice L. Update on biomarkers

for the detection of endometriosis. *Biomed Res Int* 2015;2015:130854.

- 62.** Cosar E, Mamillapalli R, Ersoy GS, Cho S, Seifer B, Taylor HS. Serum microRNAs as diagnostic markers of endometriosis: a comprehensive array-based analysis. *Fertil Steril* 2016;106:402–9.
- 63.** Bhatti Z, Salek M, Finlay A. Chronic diseases influence major life changing decisions: a new domain in quality of life research. *J R Soc Med* 2011;104:241–50.
- 64.** Apostolopoulos NV, Alexandraki KI, Gorry A, Coker A. Association between chronic pelvic pain symptoms and the presence of endometriosis. *Arch Gynecol Obstet* 2016;293:439–45.
- 65.** Droz J, Howard FM. Use of the Short-Form McGill Pain Questionnaire as a diagnostic tool in women with chronic pelvic pain. *J Minim Invasive Gynecol* 2011;18:211–7.