

Complement activation and regulation in preeclampsia and hemolysis, elevated liver enzymes, and low platelet count syndrome



Richard M. Burwick, MD, MPH; Bruce B. Feinberg, MD

The Complement System

The complement system, which comprised more than 30 soluble and membrane-bound complement proteins, is critical to human health owing to its central role in host defense and innate immunity.^{1–3} Complement proteins provide immediate defense against foreign pathogens, such as viruses and bacteria, through opsonization, inflammation, and cell membrane attack. Complement proteins are activated via enzymatic cleavage through at least 3 distinct pathways as follows: (1) the classical pathway, in which complement is activated when substances such as DNA, apoptotic cells, C-reactive protein, or immune complexes bind to complement component 1q (C1q)^{4–7}; (2) the lectin complement pathway, where complement is activated when mannose-binding lectins or ficolins bind carbohydrate patterns on foreign pathogens⁸; or (3) the alternative complement pathway, where complement is continuously activated at a low level to probe foreign, modified self-, and unal-

The complement system is critical to human health owing to its central role in host defense and innate immunity. During pregnancy, the complement system must be appropriately regulated to allow for immunologic tolerance to the developing fetus and placenta. Although some degree of complement activation can be seen in normal pregnancy, the fetus seems to be protected in part through the placental expression of complement regulatory proteins, which inhibit complement activation at different steps along the complement activation cascade. In women who develop preeclampsia and hemolysis, elevated liver enzymes, and low platelet count syndrome, there is a shift toward increased complement activation and decreased complement regulation. There is an increase in placental deposition of C5b-9, which is the terminal effector of classical, lectin, and alternative complement pathways. C5b-9 deposition stimulates trophoblasts to secrete soluble fms-like tyrosine kinase-1, which sequesters vascular endothelial growth factor and placental growth factor. Pathogenic mutations or deletions in complement regulatory genes, which predispose to increased complement activation, have been detected in women with preeclampsia and hemolysis, elevated liver enzymes, and low platelet count syndrome. Before the disease, biomarkers of alternative complement pathway activation are increased; during active disease, biomarkers of terminal complement pathway activation are increased. Urinary excretion of C5b-9 is associated with preeclampsia with severe features and distinguishes it from other hypertensive disorders of pregnancy. Taken together, existing data link preeclampsia and hemolysis, elevated liver enzymes, and low platelet count syndrome with increased activation of the terminal complement pathway that, in some cases, may be influenced by genetic alterations in complement regulators. These findings suggest that the inhibition of the terminal complement pathway, possibly through C5 blockade, may be an effective strategy to treat preeclampsia and hemolysis, elevated liver enzymes, and low platelet count syndrome, but this strategy warrants further evaluation in clinical trials.

Key words: alternative pathway, anaphylatoxins, atypical hemolytic uremic syndrome, classical pathway, complement C4, complement C5, complement inactivating agents, complement membrane attack complex, complement system proteins, HELLP syndrome, immune system, placenta, preeclampsia, pregnancy

From the Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Cedars-Sinai Medical Center, Los Angeles, CA (Dr Burwick); and Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Columbia University Irving Medical Center, New York, NY (Dr Feinberg).

Received July 14, 2020; revised Sept. 19, 2020; accepted Sept. 23, 2020.

R.M.B. has received speaking fees and research grants from Alexion Pharmaceuticals. Alexion Pharmaceuticals was not involved with this work in any manner. B.B.F. reports no conflict of interest.

This paper is part of a supplement.

Corresponding author: Richard M. Burwick, MD, MPH. richard.burwick@cshs.org

0002-9378/\$36.00

© 2020 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.ajog.2020.09.038>

tered self-structures (Figure 1).^{9,10} The indiscriminate activation of the alternative pathway is suppressed through the expression of complement regulatory proteins on healthy host surfaces.²

The complement pathways converge to activate complement protein C3 (C3), generating complement component 3a (C3a) and opsonin complement component 3b (C3b), which deposits on membrane surfaces and initiates the C3 amplification loop.¹¹ Continued deposition of C3b leads to the formation of

the complement protein C5 (C5) convertase, which activates the terminal complement pathway through cleavage of C5 to generate complement component 5a (C5a) and complement component 5b (C5b). C3a and C5a are considered anaphylatoxins because of their potent chemotactic properties and ability to increase vascular permeability and stimulate smooth muscle contractions.^{12,13} C5b combines with complement protein C6 (C6), complement protein C7 (C7), complement protein

C8 (C8), and complement protein C9 (C9) to form the terminal complement complex (C5b-9), also known as the membrane attack complex (MAC) because of its ability to form trans-membrane pores and propagate cell lysis.^{14,15} Numerous complement proteins regulate both amplification and degradation of C3b and formation of C5b-9 on the cell surface, thereby limiting harm to self.^{2,11} These complement regulators may be soluble (eg, complement factor H [CFH]—, factor I—, and factor H—related proteins) or membrane bound (eg, cluster of differentiation 55 [CD55], cluster of differentiation 46 [CD46], cluster of differentiation 55 [CD55], and cluster of differentiation 59 [CD59]).

Complement activation in pregnancy

During pregnancy, the host complement system must be appropriately regulated to allow for immunologic tolerance to the developing fetus and placenta, which are themselves semiallogenic or, in the setting of donor egg and surrogate pregnancy, fully allogenic.¹ Although some degree of complement activation can be seen in normal pregnancy^{16,17} owing to multiple factors including placental apoptotic debris, cell-free fetal DNA and RNA, and immune complex formation,^{18–22} the fetus seems to be protected in part through the placental expression of complement regulatory proteins, which inhibit complement activation at different steps along the complement activation cascade (Figure 1).^{23–25}

Various complement activation products have been measured in normal pregnancy to show that systemic complement activation is present. Richani et al¹⁶ showed that C3a and C5a are increased in healthy pregnant women at 20 to 42 weeks' gestation compared with nonpregnant women (median, C3a, 2365 vs 1340 ng/mL; C5a, 12.4 vs 4.1 ng/mL; both, $P<.001$). Derzsy et al¹⁷ showed that soluble C5b-9 levels are increased in the plasma of women with normal pregnancy at 36 to 37 weeks' gestation compared with healthy nonpregnant women (median, C5b-9, 60 vs 33 ng/mL; $P<.001$). Systemic complement

activation in pregnancy is balanced by a concomitant increase in soluble complement regulators, such as CFH,^{17,26,27} and membrane-bound complement regulators, including CD46, CD55, and CD59, which are widely expressed on placental trophoblast cells from at least 6 weeks' gestation to term.^{23–25} Endovascular trophoblasts, which invade maternal spiral arteries, also show a high expression of CD59, and this may confer additional protection against maternal complement attack.²⁸

Overall, normal pregnancy is characterized by a state of complement homeostasis, whereby an increase in systemic complement activation is balanced by a concomitant increase in complement regulatory proteins in maternal circulation and at the placental interface.

Preeclampsia and Hemolysis, Elevated Liver Enzymes, and Low Platelet Count Syndrome

Preeclampsia is defined by hypertension and proteinuria that develops in the second half of pregnancy.²⁹ When there is severe hypertension ($\geq 160/110$ mm Hg) or end-organ injury (eg, elevated serum creatinine, pulmonary edema, severe headache), with or without proteinuria, it is termed preeclampsia with severe features. Those with microangiopathic hemolysis, elevated liver enzymes, and low platelet count (HELLP) are diagnosed with HELLP syndrome.³⁰ Regardless of the terminology, pregnant women with preeclampsia without severe features, preeclampsia with severe features, or HELLP syndrome, all improve soon after the placenta is removed at delivery.

Preeclampsia has long been considered a placentally mediated disease.^{31,32} Early impairments soluble fms-like tyrosine kinase-1 (sFlt-1) in placental blood flow and oxygenation, such as with poor spiral artery remodeling,^{33,34} may increase placental inflammation and apoptosis over time, with systemic activation of leukocytes and endothelial cells.^{35,36} In women who develop preeclampsia, there is necrotic shedding of placental trophoblast into the maternal circulation³⁷ and increased trophoblast

release of sFlt-1 an antiangiogenic factor that binds and sequesters placental growth factor (PlGF) and vascular endothelial growth factor (VEGF).^{38–40} Increased levels of sFlt-1 and reduced levels of PlGF and VEGF are associated with hypertension and proteinuria and the development of preeclampsia.^{40,41}

There is an increasing recognition that complement proteins are also involved in the pathophysiology of preeclampsia and HELLP syndrome, through excess complement activation, decreased complement regulation, or both.

Placental Complement Deposition

The terminal effector of the complement pathway is C5b-9, which has both lytic and sublytic inflammatory effects. Therefore, placental deposition of C5b-9 may contribute to placental injury in women who develop preeclampsia.

In 2008, Rampersad et al⁴² showed that trophoblast deposition of C5b-9 was increased in cell culture under hypoxic conditions and C5b-9 deposition enhanced apoptotic cell death. Moreover, the surface density of C5b-9 staining and colocalization of fibrin-C5b-9 complexes were increased 2-fold in placental sections from preeclamptic pregnancies compared with those from uncomplicated pregnancies. Subsequent studies by Banadakoppa et al⁴³ and Collier et al⁴⁴ also found increased trophoblast deposition of C5b-9 in placentas from women with preeclampsia and HELLP syndrome compared with placentas from a healthy pregnancy. In the study by Collier et al,⁴⁴ the degree of placental C5b-9 staining was similar in preeclampsia and HELLP syndrome.

Although C5b-9 is the shared terminal effector of all complement pathways, investigators have sought to determine whether placental C5b-9 deposition is driven by a specific activation pathway. In 2012, Buurma et al⁴⁵ showed that placental trophoblast staining for complement component 4d (C4d) was more often detected in preeclamptic placentas than control placentas (C4d positive, 50% vs 3%; $P=.0001$).⁴⁵ The authors noted that C4d is a component of both the classical and lectin pathways, and therefore, they investigated the presence

of C1q and mannose-binding lectin to determine which pathway was responsible. Mannose-binding lectin was never observed, making it unlikely that placental C4d deposits were a result of lectin pathway activation. When C4d was present in a diffuse staining pattern at the syncytiotrophoblast surface, it colocalized with C1q, indicating that C4d deposits were most likely the result of classical pathway activation. Collier et al⁴⁴ also found increased C4d deposition in preeclamptic and HELLP placentas vs controls (Figure 2), and like their findings with C5b-9, the level of placental C4d staining was no different between preeclampsia and HELLP syndrome.⁴⁴

Although placental trophoblast deposition of C5b-9 seems to be increased in preeclamptic placentas, it remains unclear whether this is caused by increased complement activation or impaired regulation. Buurma et al⁴⁵ found that placental mRNA expressions of complement regulatory proteins CD55 and CD59 were increased 2-fold and 4-fold, respectively, in the placentas of women who delivered with preeclampsia compared with the placentas from uncomplicated deliveries. The investigators hypothesized that up-regulation of CD55 and CD59 in the face of increased complement activation suggests the presence of a feedback mechanism to maintain trophoblast integrity.


Complement and Antiangiogenic Factors

Because placental release of antiangiogenic factor sFlt-1 has been consistently implicated in the development of preeclampsia,^{40,41} there has been interest in the relationship between complement proteins and angiogenic factors. Placental ischemia and hypoxia have been shown to increase the trophoblast release of sFlt-1,^{46,47} but increasing data suggest that terminal complement proteins C5a and C5b-9 may also be implicated in this process.

In 2006, using antibody independent mouse models of complement dysregulation, Girardi et al⁴⁸ showed that complement activation products,

FIGURE 1

Schematic of complement activation and regulation at the placental interface




The complement cascade may be activated (complement activation) through distinct pathways. The classical pathway is activated by substances such as apoptotic cells, DNA, and immune complexes; the lectin pathway is activated by carbohydrate patterns on foreign surfaces; and the alternative pathway is constitutively active through spontaneous hydrolysis. These pathways converge to generate C3 convertases, which cleave C3 to generate activation products C3a (anaphylatoxin) and C3b (opsonin). The accumulation of C3b leads to the generation of C5 convertases, which cleave C5 of the terminal pathway to generate C5a (anaphylatoxin) and C5b, which combines with complement proteins C6 to C9 to form C5b-9 (MAC). C5b-9 may incorporate into cell membranes, pictured above (red cylinder on placental surface) with sublytic or lytic effects. C5b-9 may also be released in an active soluble form (soluble C5b-9), pictured above (red cylinder released from placental surface). The complement cascade may be inhibited by (1) soluble complement regulatory proteins, including CFH and CFI, which function together to inactivate C3b and down-regulate complement activation, and CFHR1, which blocks C5 convertase activity and interferes with membranous C5b-9 deposition, and (2) membrane-bound complement regulatory proteins, which may be expressed on the trophoblast membrane as pictured above (green rectangles on placental surface: CD46, CD55, CD59); CD46 blocks the activation of C3 into C3a and C3b; CD55 inhibits the actions of C3b and reduces C5 activation; and CD59 blocks the actions of C5b-9, the MAC.

C3, complement protein C3; C5, complement protein C5; C6, complement protein C6; C9, complement protein C9; CFH, complement factor H; CFH, complement factor H; CFHR, complement factor H related; CFI, complement factor I; MAC, membrane attack complex.

Buurma. Complement in preeclampsia and HELLP syndrome. Am J Obstet Gynecol 2022.

particularly C5a, induced the dysregulation of angiogenic factors. Specifically, C5a stimulated monocytes to produce soluble VEGF receptor 1 (VEGFR1) (also known as sFlt-1), which sequestered VEGF and contributed to abnormal placental development, fetal growth restriction, and death. When the effects of C5a were blocked, by inhibiting

the interaction of C5a with its receptor complement component 5a receptor (C5aR), the release of sFlt-1 was prevented and fetal death was averted. Langer et al⁴⁹ subsequently showed that C5a was capable of polarizing macrophages toward an antiangiogenic phenotype, through increased expression and secretion of soluble VEGFR1

FIGURE 2**Placental complement activity by immunofluorescence studies**

Representative C4d (green) with DAPI nuclear staining (blue) in the placenta of control, PE, and HELLP is depicted. Reproduced, with permission, from Yonekura Collier et al.⁴⁴

DAPI, 4',6-diamidino-2-phenylindole; HELLP, hemolysis, elevated liver enzymes, and low platelet count.

Burwick. Complement in preeclampsia and HELLP syndrome. *Am J Obstet Gynecol* 2022.

(sFlt-1). The supernatant from C5a-stimulated human monocytes, enriched with sFlt-1, inhibited the VEGF-induced capillary-like sprout formation of human umbilical vein endothelial cells, suggesting that proangiogenic endothelial cell functions were impaired. Normal angiogenesis was restored with antibody blockade of C5.

These initial studies showed that C5a could polarize monocytes and macrophages toward an antiangiogenic phenotype, through the secretion of sFlt-1. In 2018, Ma et al⁵⁰ showed that C5aR was also expressed on placental trophoblasts, and therefore, C5a could polarize them toward an antiangiogenic phenotype. Investigators showed that C5a stimulation of trophoblasts led to increased mRNA levels of sFlt-1 and reduced levels of PlGF, which was

further confirmed by immunofluorescent analysis.⁵⁰ C5a stimulation also reduced trophoblast migration and capillary-like tube formation, but this effect was mitigated by C5aR inhibition. Banadakoppa et al⁴³ took these experiments a step further by correlating trophoblast release of sFlt-1 with the degree of complement activation. Investigators observed increased trophoblast levels of sFlt-1 with increased complement activation, but as the degree of complement activation increased, cellular levels of sFlt-1 decreased. They hypothesized that complement activation induced secretion of sFlt-1 protein from trophoblast cells. To confirm this, they measured sFlt-1 protein levels in cell culture and they found that levels increased in conjunction with greater complement activation. Moreover, secretion of sFlt-1 correlated most closely with trophoblast release of C5b-9. In 2019, Collier et al⁴⁴ also showed that placental C5b-9 and sFlt-1 levels were strongly associated ($R=0.59$, $P=.01$) and placental sFlt-1 signaling was spatially correlated in proximity to syncytiotrophoblast membrane C4d and C5b-9 immunofluorescence signal.⁴⁴

Complement Gene Mutations

Knowledge of complement gene mutations has increased significantly over the past decade, in large part owing to a better understanding of atypical hemolytic uremic syndrome (aHUS), which is a complement-mediated disorder. Complement gene mutations are detected in approximately 60% of aHUS cases,^{51,52} including pregnancy-associated cases.^{53–55} Most genetic aHUS cases are heterozygous and are attributed to the genes CFH (25% to 30% of cases), CD46 (8% to 10%), and C3 and complement factor I (CFI) (4% to 8%, respectively).^{56,57} Mutations with a minor allele frequency of 1% to 5% have generally been referred to as uncommon variants, whereas those with a minor allele frequency of <1% are considered rare variants.⁵⁸ Determining whether a complement gene mutation is benign or pathogenic (clinically meaningful) is challenging, and such

determinations have relied on functional testing or on detailed statistical analysis of allele frequencies in patient cohorts compared with reference genomic datasets.^{56,59,60}

Complement gene mutations have also been identified in pregnant women with preeclampsia and HELLP syndrome without apparent aHUS, and these studies are presented in Table 1.^{61–65} The clinical phenotype of aHUS and HELLP syndrome overlap,^{55,66} and the 2 conditions can be difficult to distinguish in the peripartum period, although HELLP syndrome usually resolves after delivery, whereas aHUS typically does not. Nonetheless, the overlapping clinical phenotype of these conditions suggests that there may be shared biologic underpinnings. An underlying complement gene mutation could help to explain why some pregnant women develop severe preeclampsia and HELLP syndrome in the setting of complement activation whereas others do not.

Complement Mutations in Preeclampsia and Hemolysis, Elevated Liver Enzymes, and Low Platelet Count Syndrome

In 2007, Fang et al⁶⁷ reported a patient with HELLP syndrome who carried a membrane cofactor protein (MCP) gene variant (A304V) known to be associated with aHUS. The A304V variant was found in the transmembrane domain of MCP, and compared with wild type, the variant allowed more C3b deposition in response to increased antibody challenge. The authors concluded that the A304V variant in MCP had a reduced functional ability to regulate complement activation in situ. Subsequently, numerous investigators have evaluated the MCP gene in preeclampsia and HELLP syndrome. From a total of 264 women with preeclampsia or HELLP syndrome, MCP variants were detected in 8% of cases (21/264) (Table 1).^{61–65} However, the prevalence ranged widely (0%–12%), possibly caused by heterogeneous clinical phenotypes and ethnic variation.

TABLE 1
Complement gene mutations and deletions in preeclampsia and HELLP syndrome

Author, year	Case population ^a	N	Complement genes tested	Any variant ^b	C3	CFB	CFH	CFHR1, CFHR3, CFHR5	CFI	MCP
Fakhouri et al, ⁶¹ 2008	HELLP and renal involvement	11	CFH, CFI, MCP	4/11 (36)	N/A	N/A	1/11 (9.1)	N/A	2/11 (18)	1/11 (9.1)
Salmon et al, ⁶² 2011	SLE or APL Ab with preeclampsia or HELLP	40	CFH, CFI, MCP	7/40 (18)	N/A	N/A	1/40 (2.5)	N/A	2/40 (5.0)	4/40 (10)
Salmon et al, ⁶² 2011	Nonautoimmune severe preeclampsia or HELLP	59	CFH, CFI, MCP	5/59 (8.5)	N/A	N/A	0/59 (0)	N/A	1/59 (1.7)	4/59 (6.8)
Crovetto et al, ⁶³ 2012	HELLP	33	C3, CFB, CFH, CFI, MCP	2/33 (6.1)	0/33 (0)	0/33 (0)	0/33 (0)	N/A	1/33 (3.0)	1/33 (3.0)
Lokki et al, ⁶⁴ 2015	Severe preeclampsia	95	MCP	11/95 (12)	N/A	N/A	N/A	N/A	N/A	11/95 (12)
Vaught et al, ⁶⁵ 2018	Partial HELLP	14	C3, CFB, CFH, CFHR1, CFHR3, CFHR5, CFI, MCP	3/14 (21)	2/14 (14)	0/14 (0)	0/14 (0)	3/14 (21)	1/14 (7.1)	0/14 (0)
Vaught et al, ⁶⁵ 2018	HELLP	11	C3, CFB, CFH, CFHR1, CFHR3, CFHR5, CFI, MCP	5/11 (45)	1/11 (9.1)	0/11 (0)	0/11 (0)	4/11 (36)	0/11 (0)	0/11 (0)
Total	Preeclampsia or HELLP	263	C3, CFB, CFH, CFHR1, CFHR3, CFHR5, CFI, MCP	37/263 (14)	3/58 (5.2)	0/58 (0)	2/168 (1.2)	7/25 (28)	7/168 (4.2)	21/263 (8.0)

Data are presented as number/total number (percentage).

APL Ab, antiphospholipid antibody positive; C3, complement protein C3; CFB, complement factor B; CFH, complement factor H; CFHR, complement factor H related; CFI, complement factor I; HELLP, hemolysis, elevated liver enzymes, and low platelet count syndrome; MCP, membrane cofactor protein (CD46); N/A, not assessed; SLE, systemic lupus erythematosus.

^a Descriptions per investigators; severe preeclampsia and partial HELLP now termed preeclampsia with severe features; ^b Any variant of the ones listed in column 4, complement genes tested.

Burwick. Complement in preeclampsia and HELLP syndrome. *Am J Obstet Gynecol* 2022.

In a cohort of 40 women with pre-eclampsia and autoimmune disease (lupus or antiphospholipid antibodies), Salmon et al⁶² found MCP variants in 4 women (10%), 3 of whom had the A304V variant. There were no MCP variants found in 34 matched controls. Investigators then evaluated the A304V MCP variant in women with pre-eclampsia or HELLP syndrome, but without autoimmune disease, and the variant was found in 6.8% of cases (4/59) compared with 2.1% of matched controls with healthy pregnancy (3/143). However, other studies of MCP (CD46) gene variants in preeclampsia and HELLP syndrome have been less compelling. Lokki et al⁶⁴ evaluated the A304V MCP variant in 95 primiparous women with severe preeclampsia in a Finnish cohort. Heterozygosity for the A304V variant was observed in 12% of preeclampsia cases (11/95) compared with 11% of healthy controls (21/188) (odds ratio [OR], 0.94; 95% confidence interval [CI], 0.45–2.0). Notably, 3 other studies, with a combined 55 women with HELLP syndrome, detected MCP variants in only 3.6% of cases (2/55).^{61,63,65}

Of the studies listed in Table 1, mutations in CFH and CFI were detected in 1.2% (2 of 168) and 4.2% (7/168) of women with preeclampsia or HELLP syndrome, respectively. Both CFH and CFI are soluble complement regulators that function to inactivate C3b and thereby down-regulate complement activation. Pathogenic mutations in CFH and CFI are usually loss-of-function, leading to increased C3b deposition and increased complement activation. For example, Fakhouri et al⁶¹ described a patient with HELLP syndrome who had a CFH mutation (pArg303Gln) located in the SCR-1/5 domain important for C3b binding and decay-accelerating activity. Another patient with HELLP syndrome had a CFI mutation (pArg345Gln, c1034G>A), with functional analysis demonstrating a marked defect in both C3b and C4b cofactor activities.

In 2018, Vaught et al⁶⁵ analyzed an expanded number of complement genes to identify rare variants and deletions in

CFH-related genes (CFH-related protein 1 [CFHR1], CFH-related protein 3 [CFHR3], and CFH-related protein 5 [CFHR5]). Homozygous CFHR1-CFHR3 deletion has been reported in approximately 2% of the population and has a known association with aHUS.⁶⁸ Complement gene variants or CFHR gene deletions were detected in 21% of women with partial HELLP (now called preeclampsia with severe features) (3 of 14) and 45% of women with HELLP syndrome (5 of 11). Notably, 5 of these women had homozygous deletions in CFHR genes, including 4 with homozygous CFHR1-CFHR3 deletion (3 cases of preeclampsia with severe features, 1 case of HELLP) and 1 with homozygous CFHR1 deletion (HELLP). CFHR1 blocks C5 convertase activity and interferes with membranous C5b-9 deposition,⁶⁹ and therefore, deletion of CFHR1 may result in increased C5b-9 formation.

It should be noted that most complement genetic studies have assessed maternal but not fetal genetic status. The fetus has 50% chance of inheriting a heterozygous maternal variant (concordant maternal-fetal status) but 50% chance of not inheriting it (discordant maternal-fetal status). In addition, the fetus may inherit complement gene mutations from the paternal side. Thus, fetal complement genetic testing is urgently needed in future investigations.

Complement Biomarkers Before the Disease

The presence of complement genetic variants may predispose some women to develop preeclampsia and HELLP syndrome. However, even when there is a known complement gene mutation, penetrance is limited. Thus, it would be helpful to determine whether abnormal complement activation is present before the onset of disease.

In 1989, Haeger et al⁷⁰ showed that women who developed preeclampsia had higher plasma C3a levels, but not C5a levels, 1 month before delivery compared with unmatched healthy controls. Nearly 20 years later, Lynch et al⁷¹ evaluated alternative complement pathway fragment Bb in a prospective

study of 701 women. Investigators found that women with Bb levels in the upper 90th percentile before 20 weeks' gestation had 3.3 times greater relative risk of preeclampsia (adjusted OR [aOR], 3.8; 95% CI, 1.6–9; $P=.002$) after adjustment for confounders. In contrast, soluble C5b-9 levels were not increased in early gestation (10–15 weeks) among women who developed preeclampsia.⁷² Taken together, these results show that the activation of the alternative complement pathway, but not the terminal complement pathway, is increased in early pregnancy among women who develop preeclampsia.

In a larger cohort of women ($n=1002$) with plasma samples collected before 20 weeks' gestation, Lynch et al⁷³ showed that C3a levels were higher in women who developed any hypertensive disorder of pregnancy (848 vs 757 ng/L; $P<.001$), but not preeclampsia specifically. Investigators noted that the development of preeclampsia was most likely in those with underlying obesity (body mass index, >30 kg/m²) and elevated Bb levels (aOR, 10.0; 95% CI, 3.3–30; $P<.001$) or C3a levels (aOR, 8.8; 95% CI, 3–24; $P<.001$) before 20 weeks' gestation.⁷⁴ These results demonstrated a combined impact of obesity and elevated complement on the development of preeclampsia.

In 2015, Banadakoppa et al⁷⁵ evaluated complement split products in amniotic fluid collected from 731 women at the time of genetic amniocentesis. Compared with women with uncomplicated term pregnancy, those who developed early-onset preeclampsia before 34 weeks' gestation had higher amniotic fluid C3a levels (318.7 vs 254.5 ng/mL; $P=.04$) and Bb levels (1127 vs 749 ng/mL; $P=.03$). Median levels of C4a and C5a were not significantly different between the groups. These results show that complement activation is not only increased in the plasma in early pregnancy but also in the amniotic fluid before the onset of preeclampsia.

Previous studies evaluated complement markers in samples collected mostly between 10 and 20 weeks' gestation. In 2020, He et al⁷⁶ evaluated 500 women, using plasma samples collected

TABLE 2
Complement biomarkers in preeclampsia and HELLP syndrome

Author, year	Cases ^a (n)	Controls ^b (n)	Specimen	Fold change: cases vs controls		
				C3a	C5a	C5b-9
Haeger et al, ⁷⁰ 1989	Preeclampsia (14)	Healthy, unmatched (16)	Plasma	4.0×	2.6×	N/A
Haeger et al, ⁷⁷ 1990	HELLP (10)	Healthy, unmatched (10)	Plasma	5.6×	3.2×	n.d.
Derzsy et al, ¹⁷ 2010	Preeclampsia (60)	Healthy, unmatched (60)	Plasma	1.8×	N/A	1.3×
Soto et al, ⁷⁸ 2013	Preeclampsia without SGA (54)	Healthy, unmatched (134)	Plasma	n.d.	1.6×	N/A
Soto et al, ⁷⁸ 2013	Preeclampsia with SGA (52)	Healthy, unmatched (134)	Plasma	n.d.	1.6×	N/A
Burwick et al, ⁷⁹ 2013	Severe preeclampsia (25)	Healthy, GA matched (25)	Plasma	n.d.	1.3×	1.3×
Agostinis et al, ⁸⁰ 2016	Preeclampsia (30)	Healthy, GA matched (30)	Plasma	N/A	n.d.	n.d.
He et al, ⁸¹ 2016	Early-onset severe preeclampsia (30)	Healthy, GA matched (30)	Plasma	18×	2.7×	2.8×
He et al, ⁸¹ 2016	Late-onset severe preeclampsia (30)	Healthy, GA matched (30)	Plasma	115×	6.1×	2.2×
Burwick et al, ⁸² 2018	Preeclampsia with severe features (104)	Healthy, GA matched (54)	Plasma	N/A	N/A	2.0×
Burwick et al, ⁸³ 2019	Preeclampsia (16)	Healthy, unmatched (16)	Plasma	N/A	n.d.	1.4×
Burwick et al, ⁷⁹ 2013	Severe preeclampsia (25)	Healthy, GA matched (25)	Urine	3.7×	5.4×	>4.3×
Burwick et al, ⁸² 2018	Preeclampsia with severe features (104)	Healthy, GA matched (54)	Urine	N/A	N/A	4.5×
Burwick et al, ⁸³ 2019	Preeclampsia (16)	Healthy, unmatched (16)	Urine	N/A	n.d.	14×

Data are fold change in cases vs controls, with $P < .05$ unless stated otherwise.

GA, gestational age; HELLP, hemolysis, elevated liver enzymes, and low platelet count; N/A, not applicable; n.d., no difference; SGA, small for gestational age.

^a Descriptions per investigators; severe preeclampsia is now termed preeclampsia with severe features; ^b Controls labeled as unmatched if the matching criteria were not described in the study design.

Burwick. Complement in preeclampsia and HELLP syndrome. *Am J Obstet Gynecol* 2022.

even earlier at 6 to 8 weeks' gestation. Samples from 21 women who developed preeclampsia were compared with 63 samples from women who had an uncomplicated pregnancy. Investigators found that alternative pathway factors CFB and CFH were increased in women who developed preeclampsia compared with those who did not (CFB, 483 vs 403 mg/L; $P = .004$; CFH, 456 vs 366 mg/L; $P = .002$), whereas levels of C3a, C5a, and sC5b-9 were no different. These results show that even as early as 6 to 8 weeks, there is evidence of increased activation of the alternative complement pathway.

Complement Biomarkers in Active Disease

Biomarker studies suggest that before the onset of preeclampsia and HELLP syndrome, there is an early strain on the alternative complement pathway, but not the terminal pathway. However, data have shown significant activation of the terminal complement pathway in the

setting of active disease. Summary findings from complement biomarker studies among women with active preeclampsia and HELLP syndrome are presented in Table 2.^{78–83}

In 1989, Haeger et al^{70,77} first reported an increase in both C3a and C5a in women with preeclampsia and HELLP syndrome. Subsequent studies in the past decade have been mixed, in part owing to varying study design and assay systems. Plasma C3a levels were increased compared with healthy controls in 4 studies^{17,70,77,81} whereas 2 others found no difference.^{78,79} Plasma C5a levels were increased in preeclampsia or HELLP compared with healthy controls in 5 studies^{70,77–81} whereas 2 others found no difference.^{80,83} Notably, 5 studies found that plasma C5b-9 levels were increased in preeclampsia^{17,78,79,82,83} whereas 2 studies found no difference.^{77,80}

All complement biomarker studies among women with severe preeclampsia

(now preeclampsia with severe features) showed a significant increase in plasma C5a and C5b-9 levels compared with healthy controls, suggesting that terminal complement activation may be more consistently elevated in severe disease. However, Burwick et al^{79,82} showed that plasma C5a and C5b-9 levels did not differentiate preeclampsia with severe features from other hypertensive disorders of pregnancy, including gestational hypertension, chronic hypertension, and preeclampsia without severe features. These findings suggest that plasma markers of terminal complement activation (C5a and C5b-9) may be broadly increased in hypertensive disorders of pregnancy.

In urine studies, Burwick et al⁷⁹ showed that C5a levels were increased 5-fold in women with severe preeclampsia compared with matched healthy controls. Urine C5b-9 levels were also increased more than 4-fold among women with severe preeclampsia, and

this finding was seen across 3 different studies.^{79,82,83} In contrast to plasma C5b-9 levels, increased urinary C5b-9 levels were specific to women with severe preeclampsia (median [interquartile range] urine C5b-9: severe preeclampsia, 4.3 [1.2–15.1] ng/mL; chronic hypertension, 0 [0–0] ng/mL; healthy controls, 0 [0–0] ng/mL; $P<.0001$).⁷⁹ Urinary excretion of C5b-9 was detected in 96% of cases with severe preeclampsia, 12% of controls with chronic hypertension, and 8% of healthy controls.

In a multicenter observational study, Burwick et al⁸² showed that urine C5b-9 concentrations differentiated preeclampsia with severe features from other hypertensive disorders of pregnancy (chronic hypertension, gestational hypertension, or preeclampsia without severe features), with area under the receiver operating characteristic curve of 0.74 (95% CI, 0.68–0.80). Among women with preeclampsia with severe features, 40% (42 of 104) had urine C5b-9 concentration of ≥ 22 ng/mL compared with 0% of women with chronic or gestational hypertension (0 of 137) ($P<.001$) and 11% of those with preeclampsia without severe features (6 of 57) ($P<.001$). After multivariable logistic regression, preeclampsia with severe features was more likely in women with urine C5b-9 of ≥ 22 ng/mL (aOR, 10.0; 95% CI, 3.5–29; $P<.001$).

Under normal circumstances, urinary excretion of C5b-9 is not expected because of its large size ($>1,000,000$ Daltons).⁸⁴ However, C5b-9 may form at the glomerular membrane with shedding into the urine, or C5b-9 may result from complement-mediated inflammation and cellular injury at the proximal tubule.^{85–89} Burwick et al⁸⁷ showed that urinary excretion of C5b-9 increased in association with other urinary biomarkers indicative of glomerular and proximal tubule injury. Guseh et al⁹⁰ showed that in comparison with women without C5b-9 in urine, those with detectable urinary C5b-9 levels had significantly increased plasma concentrations of sFlt-1 (32,029 vs 4556 pg/mL; $P<.0001$) and significantly decreased concentrations of PlGF (15.6 vs 226 pg/

mL; $P<.0001$) and VEGF (119 vs 153 pg/mL; $P=.001$). These findings demonstrated that urinary excretion of C5b-9 correlated strongly with the antiangiogenic state in women with preeclampsia. Although these findings are intriguing, urinary C5b-9 measurements are not yet validated for clinical use, limiting immediate applicability.

Terminal Complement Blockade

Taken together, existing data show that complement regulation is impaired, and terminal complement activation is increased, in many women who develop preeclampsia and HELLP syndrome. These findings suggest that inhibition of the terminal complement pathway (eg, C5 blockade) may be an effective strategy to mitigate disease.

The only C5 blocker currently used in pregnant and lactating women is eculizumab, a humanized monoclonal antibody against C5, which inhibits cleavage of C5 into C5a and C5b and prevents formation of the terminal complement complex C5b-9.⁹¹ It was approved by the Food and Drug Administration to treat paroxysmal nocturnal hemoglobinuria (PNH) in 2007 and aHUS in 2011, owing to its therapeutic efficacy in both conditions.^{92,93} PNH is an acquired hemolytic disorder that occurs when hematopoietic stem cells in the bone marrow lose their ability to anchor complement regulatory proteins CD55 and CD59 to the cell surface, thereby predisposing to complement-mediated red cell hemolysis. In contrast, aHUS is a complement-mediated thrombotic microangiopathy disorder, often attributed to inherited complement mutations.^{51,52} However, the penetrance of complement gene mutations is limited, approximately 50% by the age of 45 years,⁹⁴ and aHUS may not manifest clinically until there is a complement amplifying event, such as pregnancy.^{53,55}


Pregnancy safety data for eculizumab stem from its use in pregnant women with PNH and aHUS.^{55,95} Among pregnant women with PNH treated with eculizumab, Kelly et al⁹⁵ found that eculizumab was detectable in 7 of 20 cord blood samples (range, 11.8–21.1 ug/mL), suggesting that it crosses the

placenta at low levels. Although immunoglobulin G monoclonal antibodies are transported across the placenta through the neonatal crystallizable fragment receptor (FcRn), eculizumab was engineered with human immunoglobulin G2 and immunoglobulin G4 heavy-chain sequences to form a hybrid constant region with reduced affinity for FcRn.^{91,96} Hallstensen et al⁹⁷ showed that treatment with eculizumab during pregnancy does not seem to alter the complement system activity of the newborn. Kelly et al⁹⁵ also examined 10 breast milk samples among women receiving eculizumab, and the drug was not detected in any of the samples, leading the authors to conclude that breastfeeding is safe. However, continued vigilance is warranted given the small sample size.

Complement Blockade in Preeclampsia and Hemolysis, Elevated Liver Enzymes, and Low Platelet Count Syndrome

Data from PNH and aHUS have been used to strengthen the argument for C5 blockade in preeclampsia and HELLP syndrome. In ex vivo experiments, Vaught et al⁹⁸ showed that complement-mediated hemolysis was increased when the serum of women with aHUS, severe preeclampsia, or HELLP syndrome was mixed with PNH red cells lacking CD55 and CD59 (modified Ham test). Likewise, Palomo et al⁹⁹ showed that C5b-9 deposition on endothelial cells was increased after exposure to activated plasma from women with aHUS, severe preeclampsia, or HELLP syndrome. The magnitude of complement-mediated hemolysis and endothelial C5b-9 deposition in these 2 studies, respectively, was similar in subjects with preeclampsia, HELLP, or aHUS, suggesting that aberrant complement activation was a common feature of these disorders. Importantly, both studies showed that complement-mediated effects were mitigated by C5 blockade with eculizumab.^{98,99}

Although terminal complement blockade with eculizumab is effective for the treatment of PNH and aHUS, there is only a single case report describing the successful off-label use of eculizumab to

FIGURE 3**Laboratory measurements before and after complement blockade with eculizumab**

The figure depicts laboratory measures before and after eculizumab in a patient with preeclampsia and HELLP syndrome. **A**, Platelet count and lactate dehydrogenase measurements. **B**, Aspartate and alanine transaminase measurements. **C**, Plasma and urine C5b-9 measurements. Eculizumab was administered at hospital days 2, 9, and 15, with delivery at hospital day 19. Adapted from Burwick and Feinberg¹⁰⁰; and Burwick et al.¹⁰³

Ecu, eculizumab; HELLP, hemolysis, elevated liver enzymes, and low platelet count.

Burwick. Complement in preeclampsia and HELLP syndrome. *Am J Obstet Gynecol* 2022.

treat preeclampsia and HELLP syndrome in pregnancy.¹⁰⁰ However, others have shown that on-label use of eculizumab is effective in treating pregnant women with aHUS who develop preeclampsia or those who develop postpartum aHUS after a delivery complicated by preeclampsia and HELLP syndrome.^{55,101,102} In 2019, Lu et al¹⁰¹ described a pregnant woman who was diagnosed with aHUS at 22 weeks' gestation and initiated on eculizumab treatment. At 24 weeks' gestation, she was also diagnosed with superimposed preeclampsia and eculizumab treatment was continued, with prolongation of pregnancy by an additional 25 days. In retrospective laboratory analyses, the authors found that the sFlt-1 to PlGF ratio decreased after the initiation of eculizumab, and they hypothesized that eculizumab may have moderated the antiangiogenic activity in preeclampsia.¹⁰¹ In 2020, Lokki et al¹⁰² described a patient who developed postpartum aHUS after delivery for preeclampsia and HELLP syndrome at 34 weeks' gestation. The patient developed severe renal failure requiring dialysis, but she quickly recovered after the diagnosis of aHUS was made and treatment with eculizumab was initiated.

In 2013, Burwick and Feinberg¹⁰⁰ were the first to show that C5 blockade with eculizumab could also be used effectively to treat preeclampsia and HELLP syndrome, in the absence of aHUS. They described a previously healthy pregnant woman who developed severe preeclampsia and HELLP syndrome at 26 weeks' gestation. After the off-label use of eculizumab, there was resolution of hemolysis and thrombocytopenia and normalization of liver enzymes (Figure 3). Maternal and fetal status improved sufficiently to prolong pregnancy an additional 17 days, allowing more time for fetal maturation and decreasing the likelihood of neonatal morbidity. In a subsequent report, the authors reported that the concentration of C5b-9 in the blood and urine decreased in conjunction with C5 blockade and disease remission.¹⁰³ Although there was a good clinical outcome, the authors noted a persistent elevation of C5a after C5 blockade, and they cautioned that extrinsic factors such as thrombin may also generate C5a independent of the C5 convertase.¹⁰⁴ Finally, they also cautioned that eculizumab or eculizumab-C5 complexes may cross the placenta or deposit in the

maternal kidney, and long-term effects are unknown.^{97,105,106} Therefore, the benefits of eculizumab or other C5 monoclonal antibodies for the treatment of preeclampsia and HELLP syndrome require further evaluation in the setting of a clinical trial.

Discussion

The current literature demonstrates that complement activation and regulation are altered in women who develop preeclampsia and HELLP syndrome. The initial stages of disease may begin in early pregnancy, with complement proteins altering trophoblast development and mediating placental inflammation and apoptosis. Early activation of the alternative complement pathway and underlying medical conditions such as obesity magnify the likelihood of developing preeclampsia. Pathogenic complement gene mutations or deletions also confer an increased risk, owing to an impaired ability to regulate complement activation in pregnancy. Terminal complement pathway activation leads to the formation of C5a and C5b-9, which can polarize monocytes, macrophages, and trophoblast cells to the antiangiogenic phenotype, with the release of sFlt-1 into

the maternal circulation. In active disease, there is an increased formation of C5b-9 in the blood and urine, and urinary detection of C5b-9 differentiates preeclampsia with severe features from other hypertensive disorders of pregnancy.

Terminal complement pathway activation seems to exceed the complement regulation in women who develop preeclampsia and HELLP syndrome. Research to date suggests that the degree of complement activation is largely similar in preeclampsia with severe features and HELLP syndrome, consistent with our understanding of the disease spectrum. We found no evidence that HELLP syndrome is a distinct entity from preeclampsia, but rather the 2 conditions share pathophysiologic underpinnings, with excess complement activation as a common feature. Therefore, research in this area should evaluate women with both preeclampsia and HELLP syndrome to capture more pregnancies with complement-mediated disease.

The role of complement proteins in preeclampsia and HELLP syndrome is now evident but more studies are needed to understand how complement protein biomarkers and complement genetic variants may be used clinically to guide prediction, diagnosis, and treatment of disease. Some data may be extrapolated from existing studies on aHUS and other thrombotic microangiopathy disorders, but pregnancy is a unique time with distinct considerations for the mother and the developing fetus. Future clinical trials aimed at preventing or treating preeclampsia and HELLP syndrome should consider complement blocking therapies, including C5 blockade, but there remains an urgent and unmet need to evaluate the safety and efficacy of novel complement therapeutic agents in pregnant and lactating women. ■

ACKNOWLEDGMENTS

The authors thank Dr Nicholas Burwick for the critical review of the article (institutional affiliations: Department of Hospital and Specialty Medicine, Veterans Affairs Puget Sound Health Care System, Seattle, WA; and Division of Hematology, Department of Medicine, University of Washington School of Medicine, Seattle, WA).

GLOSSARY

aHUS: Atypical hemolytic uremic syndrome.
 Anaphylatoxin: A complement component which causes smooth muscle contraction, vasodilation, histamine release from mast cells, and enhanced vascular permeability.
 CD46: Cluster of differentiation 46, also known as membrane cofactor protein.
 CD55: Cluster of differentiation 55, also known as decay-accelerating factor.
 CD59: Cluster of differentiation 59, also known as protectin.
 CFB: Complement factor B.
 CFH: Complement factor H.
 CFHR: Complement factor H related.
 CFI: Complement factor I.
 C1q: Complement component 1q.
 C3: Complement component 3.
 C3a: Complement component 3a.
 C3b: Complement component 3b.
 C4: Complement component 4.
 C4d: Complement component 4d.
 C5: Complement component 5.
 C5a: Complement component 5a.
 C5aR: Complement component 5a receptor.
 C5b: Complement component 5b.
 C5b-9: Terminal complement complex, also known as membrane attack complex.
 HELLP: Hemolysis, elevated liver enzymes, low platelet count.
 MAC: Membrane attack complex (or C5b-9).
 MCP: Membrane cofactor protein (or CD46).
 Opsonin: A complement component which binds to foreign pathogens or cells making them more susceptible to phagocytosis.
 PIGF: Placental growth factor.
 PNH: Paroxysmal nocturnal hemoglobinuria.
 sFit-1: Soluble fms-like tyrosine kinase-1.
 VEGF: Vascular endothelial growth factor.
 VEGFR1: Vascular endothelial growth factor receptor 1.

REFERENCES

- Regal JF, Gilbert JS, Burwick RM. The complement system and adverse pregnancy outcomes. *Mol Immunol* 2015;67:56–70.
- Zipfel PF, Skerka C. Complement regulators and inhibitory proteins. *Nat Rev Immunol* 2009;9:729–40.
- Reis ES, Mastellos DC, Hajishengallis G, Lambris JD. New insights into the immune functions of complement. *Nat Rev Immunol* 2019;19:503–16.
- Claus DR, Siegel J, Petras K, Skor D, Osmand AP, Gewurz H. Complement activation by interaction of polyanions and polycations. III. Complement activation by interaction of multiple polyanions and polycations is the presence of C-reactive protein. *J Immunol* 1977;118:83–7.
- Garlatti V, Chouquet A, Lunardi T, et al. Cutting edge: C1q binds deoxyribose and heparan sulfate through neighboring sites of its recognition domain. *J Immunol* 2010;185:808–12.
- Jiang HX, Siegel JN, Gewurz H. Binding and complement activation by C-reactive protein via the collagen-like region of C1q and inhibition of these reactions by monoclonal antibodies to C-reactive protein and C1q. *J Immunol* 1991;146:2324–30.
- Nauta AJ, Trouw LA, Daha MR, et al. Direct binding of C1q to apoptotic cells and cell blebs induces complement activation. *Eur J Immunol* 2002;32:1726–36.
- Matsushita M, Endo Y, Fujita T. Cutting edge: complement-activating complex of ficolin and mannose-binding lectin-associated serine protease. *J Immunol* 2000;164:2281–4.
- Schmidt CQ, Lambris JD, Ricklin D. Protection of host cells by complement regulators. *Immunol Rev* 2016;274:152–71.
- Lachmann PJ, Nicol P. Reaction mechanism of the alternative pathway of complement fixation. *Lancet* 1973;1:465–7.
- Lachmann PJ. The amplification loop of the complement pathways. *Adv Immunol* 2009;104:115–49.
- Cochrane CG, Müller-Eberhard HJ. The derivation of two distinct anaphylatoxin activities from the third and fifth components of human complement. *J Exp Med* 1968;127:371–86.

13. Gorski JP, Hugli TE, Müller-Eberhard HJ. C4a: the third anaphylatoxin of the human complement system. *Proc Natl Acad Sci U S A* 1979;76:5299–302.
14. Mayer MM. Mechanism of cytolysis by complement. *Proc Natl Acad Sci U S A* 1972;69:2954–8.
15. Hammer CH, Nicholson A, Mayer MM. On the mechanism of cytolysis by complement: evidence on insertion of C5b and C7 subunits of the C5b,6,7 complex into phospholipid bilayers of erythrocyte membranes. *Proc Natl Acad Sci U S A* 1975;72:5076–80.
16. Richani K, Soto E, Romero R, et al. Normal pregnancy is characterized by systemic activation of the complement system. *J Matern Fetal Neonatal Med* 2005;17:239–45.
17. Derzsy Z, Prohászka Z, Rigó J Jr, Füst G, Molvarec A. Activation of the complement system in normal pregnancy and preeclampsia. *Mol Immunol* 2010;47:1500–6.
18. Feinberg BB. Preeclampsia: the death of Goliath. *Am J Reprod Immunol* 2006;55:84–98.
19. Smith SC, Baker PN, Symonds EM. Placental apoptosis in normal human pregnancy. *Am J Obstet Gynecol* 1997;177:57–65.
20. Lo YM, Corbetta N, Chamberlain PF, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997;350:485–7.
21. Poon LL, Leung TN, Lau TK, Lo YM. Circulating fetal RNA in maternal plasma. *Ann N Y Acad Sci* 2001;945:207–10.
22. Masson PL, Delire M, Cambiaso CL. Circulating immune complexes in normal human pregnancy. *Nature* 1977;266:542–3.
23. Holmes CH, Simpson KL, Okada H, et al. Complement regulatory proteins at the fetomaternal interface during human placental development: distribution of CD59 by comparison with membrane cofactor protein (CD46) and decay accelerating factor (CD55). *Eur J Immunol* 1992;22:1579–85.
24. Holmes CH, Simpson KL, Wainwright SD, et al. Preferential expression of the complement regulatory protein decay accelerating factor at the fetomaternal interface during human pregnancy. *J Immunol* 1990;144:3099–105.
25. Hsi BL, Hunt JS, Atkinson JP. Differential expression of complement regulatory proteins on subpopulations of human trophoblast cells. *J Reprod Immunol* 1991;19:209–23.
26. He YD, Xu BN, Song D, et al. Normal range of complement components during pregnancy: a prospective study. *Am J Reprod Immunol* 2020;83:e13202.
27. Lokki AI, Heikkinen-Eloranta J, Jarva H, et al. Complement activation and regulation in preeclamptic placenta. *Front Immunol* 2014;5:312.
28. Ueda M, Sato Y, Horie A, et al. Endovascular trophoblast expresses CD59 to evade complement-dependent cytotoxicity. *Mol Cell Endocrinol* 2019;490:57–67.
29. American College of Obstetricians and Gynecologists' Committee on Practice Bulletins—Obstetrics. Gestational hypertension and preeclampsia: ACOG Practice Bulletin, Number 222. *Obstet Gynecol* 2020;135:e237–60.
30. Weinstein L. Syndrome of hemolysis, elevated liver enzymes, and low platelet count: a severe consequence of hypertension in pregnancy. *Am J Obstet Gynecol* 1982;142:159–67.
31. Nesbitt RE Jr. Pathology of the placenta in toxemia. *Clin Obstet Gynecol* 1958;1:349–57.
32. Jeffcoate TN, Scott JS. Some observations on the placental factor in pregnancy toxemia. *Am J Obstet Gynecol* 1959;77:475–89.
33. Zhou Y, Damsky CH, Chiu K, Roberts JM, Fisher SJ. Preeclampsia is associated with abnormal expression of adhesion molecules by invasive cytotrophoblasts. *J Clin Invest* 1993;91:950–60.
34. Genbacev O, Joslin R, Damsky CH, Polliotti BM, Fisher SJ. Hypoxia alters early gestation human cytotrophoblast differentiation/invasion in vitro and models the placental defects that occur in preeclampsia. *J Clin Invest* 1996;97:540–50.
35. Germain SJ, Sacks GP, Sooranna SR, Sargent IL, Redman CW. Systemic inflammatory priming in normal pregnancy and preeclampsia: the role of circulating syncytiotrophoblast microparticles. *J Immunol* 2007;178:5949–56.
36. Roberts JM, Taylor RN, Musci TJ, Rodgers GM, Hubel CA, McLaughlin MK. Preeclampsia: an endothelial cell disorder. *Am J Obstet Gynecol* 1989;161:1200–4.
37. Huppertz B, Kingdom J, Caniggia I, et al. Hypoxia favours necrotic versus apoptotic shedding of placental syncytiotrophoblast into the maternal circulation. *Placenta* 2003;24:181–90.
38. Kendall RL, Thomas KA. Inhibition of vascular endothelial cell growth factor activity by an endogenously encoded soluble receptor. *Proc Natl Acad Sci U S A* 1993;90:10705–9.
39. Kendall RL, Wang G, DiSalvo J, Thomas KA. Specificity of vascular endothelial cell growth factor receptor ligand binding domains. *Biochem Biophys Res Commun* 1994;201:326–30.
40. Maynard SE, Min JY, Merchan J, et al. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin Invest* 2003;111:649–58.
41. Levine RJ, Maynard SE, Qian C, et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med* 2004;350:672–83.
42. Rampersad R, Barton A, Sadovsky Y, Nelson DM. The C5b-9 membrane attack complex of complement activation localizes to villous trophoblast injury in vivo and modulates human trophoblast function in vitro. *Placenta* 2008;29:855–61.
43. Banadakoppa M, Balakrishnan M, Yallampalli C. Upregulation and release of soluble fms-like tyrosine kinase receptor 1 mediated by complement activation in human syncytiotrophoblast cells. *Am J Reprod Immunol* 2018;80:e13033.
44. Yonekura Collier AR, Zsengeller Z, Pernicone E, Salahuddin S, Khankin EV, Karumanchi SA. Placental sFLT1 is associated with complement activation and syncytiotrophoblast damage in preeclampsia. *Hypertens Preg* 2019;38:193–9.
45. Buurma A, Cohen D, Veraar K, et al. Preeclampsia is characterized by placental complement dysregulation. *Hypertension* 2012;60:1332–7.
46. Li H, Gu B, Zhang Y, Lewis DF, Wang Y. Hypoxia-induced increase in soluble Flt-1 production correlates with enhanced oxidative stress in trophoblast cells from the human placenta. *Placenta* 2005;26:210–7.
47. Nagamatsu T, Fujii T, Kusumi M, et al. Cytotrophoblasts up-regulate soluble fms-like tyrosine kinase-1 expression under reduced oxygen: an implication for the placental vascular development and the pathophysiology of preeclampsia. *Endocrinology* 2004;145:4838–45.
48. Girardi G, Yarinlin D, Thurman JM, Holers VM, Salmon JE. Complement activation induces dysregulation of angiogenic factors and causes fetal rejection and growth restriction. *J Exp Med* 2006;203:2165–75.
49. Langer HF, Chung KJ, Orlova VV, et al. Complement-mediated inhibition of neovascularization reveals a point of convergence between innate immunity and angiogenesis. *Blood* 2010;116:4395–403.
50. Ma Y, Kong LR, Ge Q, et al. Complement 5a-mediated trophoblasts dysfunction is involved in the development of preeclampsia. *J Cell Mol Med* 2018;22:1034–46.
51. Kavanagh D, Richards A, Atkinson J. Complement regulatory genes and hemolytic uremic syndromes. *Annu Rev Med* 2008;59:293–309.
52. Fremeaux-Bacchi V, Fakhouri F, Garnier A, et al. Genetics and outcome of atypical hemolytic uremic syndrome: a nationwide French series comparing children and adults. *Clin J Am Soc Nephrol* 2013;8:554–62.
53. Bruel A, Kavanagh D, Noris M, et al. Hemolytic uremic syndrome in pregnancy and postpartum. *Clin J Am Soc Nephrol* 2017;12:1237–47.
54. Fakhouri F, Roumenina L, Provot F, et al. Pregnancy-associated hemolytic uremic syndrome revisited in the era of complement gene mutations. *J Am Soc Nephrol* 2010;21:859–67.
55. Gupta M, Govindappagari S, Burwick RM. Pregnancy-associated atypical hemolytic uremic syndrome: a systematic review. *Obstet Gynecol* 2020;135:46–58.
56. Osborne AJ, Breno M, Borsa NG, et al. Statistical validation of rare complement variants provides insights into the molecular basis of atypical hemolytic uremic syndrome and C3 glomerulopathy. *J Immunol* 2018;200:2464–78.
57. Nester CM, Barbour T, de Cordoba SR, et al. Atypical aHUS: state of the art. *Mol Immunol* 2015;67:31–42.
58. Manolio TA, Collins FS, Cox NJ, et al. Finding the missing heritability of complex diseases. *Nature* 2009;461:747–53.
59. Duzkale H, Shen J, McLaughlin H, et al. A systematic approach to assessing the clinical significance of genetic variants. *Clin Genet* 2013;84:453–63.

60. Walsh R, Thomson KL, Ware JS, et al. Reassessment of Mendelian gene pathogenicity using 7,855 cardiomyopathy cases and 60,706 reference samples. *Genet Med* 2017;19:192–203.
61. Fakhouri F, Jablonski M, Lepercq J, et al. Factor H, membrane cofactor protein, and factor I mutations in patients with hemolysis, elevated liver enzymes, and low platelet count syndrome. *Blood* 2008;112:4542–5.
62. Salmon JE, Heuser C, Triebwasser M, et al. Mutations in complement regulatory proteins predispose to preeclampsia: a genetic analysis of the PROMISSE cohort. *PLoS Med* 2011;8:e1001013.
63. Crovetto F, Borsa N, Acaia B, et al. The genetics of the alternative pathway of complement in the pathogenesis of HELLP syndrome. *J Matern Fetal Neonatal Med* 2012;25:2322–5.
64. Lokki AI, Aalto-Viljakainen T, Meri S, Laivuori H; FINNPEC. Genetic analysis of membrane cofactor protein (CD46) of the complement system in women with and without preeclamptic pregnancies. *PLoS One* 2015;10:e0117840.
65. Vaught AJ, Braunstein EM, Jasem J, et al. Germline mutations in the alternative pathway of complement predispose to HELLP syndrome. *JCI Insight* 2018;3:e99128.
66. Gupta M, Feinberg BB, Burwick RM. Thrombotic microangiopathies of pregnancy: differential diagnosis. *Pregnancy Hypertens* 2018;12:29–34.
67. Fang CJ, Fremaux-Bacchi V, Liszewski MK, et al. Membrane cofactor protein mutations in atypical hemolytic uremic syndrome (aHUS), fatal Stx-HUS, C3 glomerulonephritis, and the HELLP syndrome. *Blood* 2008;111:624–32.
68. Zipfel PF, Edey M, Heinen S, et al. Deletion of complement factor H-related genes CFHR1 and CFHR3 is associated with atypical hemolytic uremic syndrome. *PLoS Genet* 2007;3:e41.
69. Heinen S, Hartmann A, Lauer N, et al. Factor H-related protein 1 (CFHR-1) inhibits complement C5 convertase activity and terminal complex formation. *Blood* 2009;114:2439–47.
70. Haeger M, Bengtson A, Karlsson K, Heideman M. Complement activation and anaphylatoxin (C3a and C5a) formation in preeclampsia and by amniotic fluid. *Obstet Gynecol* 1989;73:551–6.
71. Lynch AM, Murphy JR, Byers T, et al. Alternative complement pathway activation fragment Bb in early pregnancy as a predictor of preeclampsia. *Am J Obstet Gynecol* 2008;198:385.e1–9.
72. Lynch AM, Murphy JR, Gibbs RS, et al. The interrelationship of complement-activation fragments and angiogenesis-related factors in early pregnancy and their association with preeclampsia. *BJOG* 2010;117:456–62.
73. Lynch AM, Gibbs RS, Murphy JR, Giclas PC, Salmon JE, Holers VM. Early elevations of the complement activation fragment C3a and adverse pregnancy outcomes. *Obstet Gynecol* 2011;117:75–83.
74. Lynch AM, Eckel RH, Murphy JR, et al. Prepregnancy obesity and complement system activation in early pregnancy and the subsequent development of preeclampsia. *Am J Obstet Gynecol* 2012;206:428.e1–8.
75. Banadakoppa M, Vidaeff AC, Yallampalli U, Ramin SM, Belfort MA, Yallampalli C. Complement split products in amniotic fluid in pregnancies subsequently developing early-onset preeclampsia. *Dis Markers* 2015;2015:263109.
76. He YD, Xu BN, Wang ML, et al. Dysregulation of complement system during pregnancy in patients with preeclampsia: a prospective study. *Mol Immunol* 2020;122:69–79.
77. Haeger M, Unander M, Bengtsson A. Enhanced anaphylatoxin and terminal C5b-9 complement complex formation in patients with the syndrome of hemolysis, elevated liver enzymes, and low platelet count. *Obstet Gynecol* 1990;76:698–702.
78. Soto E, Romero R, Richani K, et al. Preeclampsia and pregnancies with small-for-gestational age neonates have different profiles of complement split products. *J Matern Fetal Neonatal Med* 2010;23:646–57.
79. Burwick RM, Fichorova RN, Dawood HY, Yamamoto HS, Feinberg BB. Urinary excretion of C5b-9 in severe preeclampsia: tipping the balance of complement activation in pregnancy. *Hypertension* 2013;62:1040–5.
80. Agostinis C, Stampalija T, Tannetta D, et al. Complement component C1q as potential diagnostic but not predictive marker of preeclampsia. *Am J Reprod Immunol* 2016;76:475–81.
81. He Y, Xu B, Song D, Yu F, Chen Q, Zhao M. Expression of the complement system's activation factors in plasma of patients with early/late-onset severe pre-eclampsia. *Am J Reprod Immunol* 2016;76:205–11.
82. Burwick RM, Velásquez JA, Valencia CM, et al. Terminal complement activation in preeclampsia. *Obstet Gynecol* 2018;132:1477–85.
83. Burwick RM, Togioka BM, Speranza RJ, et al. Assessment of blood-brain barrier integrity and neuroinflammation in preeclampsia. *Am J Obstet Gynecol* 2019;221:269.e1–8.
84. Bhakdi S, Tranum-Jensen J. Molecular weight of the membrane C5b-9 complex of human complement: characterization of the terminal complex as a C5b-9 monomer. *Proc Natl Acad Sci U S A* 1981;78:1818–22.
85. Peng Q, Li K, Smyth LA, et al. C3a and C5a promote renal ischemia-reperfusion injury. *J Am Soc Nephrol* 2012;23:1474–85.
86. Zhou W, Farrar CA, Abe K, et al. Predominant role for C5b-9 in renal ischemia/reperfusion injury. *J Clin Invest* 2000;105:1363–71.
87. Burwick RM, Easter SR, Dawood HY, Yamamoto HS, Fichorova RN, Feinberg BB. Complement activation and kidney injury molecule-1-associated proximal tubule injury in severe preeclampsia. *Hypertension* 2014;64:833–8.
88. Kerjaschki D, Schulze M, Binder S, et al. Transcellular transport and membrane insertion of the C5b-9 membrane attack complex of complement by glomerular epithelial cells in experimental membranous nephropathy. *J Immunol* 1989;143:546–52.
89. Morita Y, Ikeguchi H, Nakamura J, Hotta N, Yuzawa Y, Matsuo S. Complement activation products in the urine from proteinuric patients. *J Am Soc Nephrol* 2000;11:700–7.
90. Guseh SH, Feinberg BB, Dawood HY, Yamamoto HS, Fichorova RN, Burwick RM. Urinary excretion of C5b-9 is associated with the anti-angiogenic state in severe preeclampsia. *Am J Reprod Immunol* 2015;73:437–44.
91. Thomas TC, Rollins SA, Rother RP, et al. Inhibition of complement activity by humanized anti-C5 antibody and single-chain Fv. *Mol Immunol* 1996;33:1389–401.
92. Hillmen P, Young NS, Schubert J, et al. The complement inhibitor eculizumab in paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 2006;355:1233–43.
93. Legendre CM, Licht C, Muus P, et al. Terminal complement inhibitor eculizumab in atypical hemolytic-uremic syndrome. *N Engl J Med* 2013;368:2169–81.
94. Sullivan M, Eric Z, Hoffmann MM, et al. Epidemiological approach to identifying genetic predispositions for atypical hemolytic uremic syndrome. *Ann Hum Genet* 2010;74:17–26.
95. Kelly RJ, Höchsmann B, Szer J, et al. Eculizumab in pregnant patients with paroxysmal nocturnal hemoglobinuria. *N Engl J Med* 2015;373:1032–9.
96. Brodsky RA. How I treat paroxysmal nocturnal hemoglobinuria. *Blood* 2009;113:6522–7.
97. Hallstensen RF, Bergseth G, Foss S, et al. Eculizumab treatment during pregnancy does not affect the complement system activity of the newborn. *Immunobiology* 2015;220:452–9.
98. Vaught AJ, Gavrilaki E, Hueppchen N, et al. Direct evidence of complement activation in HELLP syndrome: a link to atypical hemolytic uremic syndrome. *Exp Hematol* 2016;44:390–8.
99. Palomo M, Blasco M, Molina P, et al. Complement activation and thrombotic microangiopathies. *Clin J Am Soc Nephrol* 2019;14:1719–32.
100. Burwick RM, Feinberg BB. Eculizumab for the treatment of preeclampsia/HELLP syndrome. *Placenta* 2013;34:201–3.
101. Lu AB, Lazarus B, Rolnik DL, Palmer KR. Pregnancy prolongation after eculizumab use in early-onset preeclampsia. *Obstet Gynecol* 2019;134:1215–8.
102. Lokki AI, Haapio M, Heikkinen-Eloranta J. Eculizumab treatment for postpartum HELLP syndrome and aHUS-case report. *Front Immunol* 2020;11:548.
103. Burwick RM, Burwick NR, Feinberg BB. Eculizumab fails to inhibit generation of C5a in vivo. *Blood* 2014;124:3502–3.
104. Krisinger MJ, Goebeler V, Lu Z, et al. Thrombin generates previously unidentified C5 products that support the terminal complement activation pathway. *Blood* 2012;120:1717–25.
105. Burwick RM, Burwick N, Feinberg BB. Response: maternal and cord C5a in response to eculizumab. *Blood* 2015;126:279–80.
106. Herlitz LC, Bombach AS, Markowitz GS, et al. Pathology after eculizumab in dense deposit disease and C3 GN. *J Am Soc Nephrol* 2012;23:1229–37.